

10/28/97


**TOWNSEND and TOWNSEND and CREW LLP**

Two Embarcadero Center, 8th Floor  
San Francisco, CA 94111-3834  
(415) 576-0200

**RULE 60**

 Atty. Docket No. 16930-000921

 "Express Mail" Label No. EM197112005US

 Date of Deposit October 28, 1997

**ASSISTANT COMMISSIONER FOR PATENTS**  
**BOX PATENT APPLICATION**  
**Washington, D.C. 20231**

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231

Sir:

This is a request under 37 CFR 1.60 for filing a

☐ Continuation ☒ Division

 of application No. 08/328,673, filed October 25, 1994,

 of (list each inventor) Richard J. Gregory, Ken N. Wills, Daniel C. Maneval

By

John P. Borg

 for RECOMBINANT ADENOVIRAL VECTOR AND METHODS OF USE

The application papers **FILED HEREWITH** (specification, claims, originally filed drawing(s) and oath or declaration) are a true copy of the prior application.

☒ Please amend the specification by inserting before the first line the sentence:

--This is a ☐ Continuation ☒ Division of application No. 08/328,673 filed October 25, 1994, which is a continuation-in-part of USSN 08/233,777, filed April 26, 1994 (now abandoned), which is a continuation-in-part of USSN 08/142,669, filed October 25, 1993 (now abandoned), the disclosures of which are incorporated by reference in their entirety.--

☒ A preliminary amendment is enclosed.

☐ Formal drawings are enclosed.

☒ An Information Disclosure Statement under 37 CFR 1.97 is enclosed.

☐ Enclosed is a petition to extend time to respond.

☐ Please record the enclosed assignment to \_\_\_\_\_

☒ The prior application is assigned to Canji, Inc. (Reel 7591, Frames 0355-0359)
☒ Please cancel claim(s) 1-15, 25
☒ Copy of Revocation and Substitution of Power of Attorney

☒ Sequence Listing, Communication regarding computer readable sequence listing

**Claims as Filed, Less any Cancelled Claims**
**OTHER THAN A**
**SMALL ENTITY**

	(Col. 1)	(Col. 2)
FOR:	NO. FILED	NO. EXTRA
BASIC FEE		
TOTAL CLAIMS	15 -20=	0
INDEP CLAIMS	3 -3=	0
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIM PRESENTED		

**SMALL ENTITY**

RATE	FEE
	\$395
x11=	\$
x40=	\$
+130=	\$
TOTAL	\$

OR

OR

OR

OR

OR

OR

RATE	FEE
	\$790
x22=	\$
x80=	\$
+260=	\$
TOTAL	\$790

\* If the difference in Col. 1 is less than zero, enter "0" in Col. 2

Please charge Deposit Account No. 20-1430 as follows:

☒ Filing fee

☒ Any additional fees associated with this paper or during the pendency of this application

☐ The issue fee set in 37 CFR 1.18 at or before mailing of the Notice of Allowance, pursuant to 37 CFR 1.311(b).

 \$ 790.00
☐ A check for \$ \_\_\_\_\_ is enclosed.

☐ 2 extra copies of this sheet are enclosed.

Respectfully submitted,

TOWNSEND and TOWNSEND and CREW LLP

Renée A. Fitts

Reg. No.: 35,136

Attorneys for Applicants

Telephone: (415) 576-0200

RULE60.TRN 9/96

A P P L I C A T I O N

for

UNITED STATES LETTERS PATENT

on

RECOMBINANT ADENOVIRAL VECTOR  
AND METHODS OF USE

by

Richard J. Gregory

Ken N. Wills

Daniel C. Maneval

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" MAILING LABEL NUMBER  
DATE OF DEPOSIT 10/25/94

POSTAL SERVICE "EXPRESS MAIL" POST OFFICE TO  
ADDITIONAL CHARGE UNDER 37 CFR 1.10 ON THE DATE INDICATED  
ADDRESSES TO THE COMMISSIONER OF PATENTS AND  
TRADEMARKS, WASHINGTON, D.C. 20540

(TYPED OR PRINTED NAME OF PERSON MAILING PAPER OR FEE)  
SIGNATURE OF PERSON MAILING PAPER OR FEE

Number of Drawings: 16

Docket No.: P-CJ 1192

Attorneys

CAMPBELL AND FLORES  
4370 La Jolla Village Drive, Suite 700  
San Diego, California 92122

RECOMBINANT ADENOVIRAL VECTOR AND METHODS OF USE  
BACKGROUND OF THE INVENTION

This application is a continuation-in-part of U.S. Serial No. 08/233,777, filed May 19, 1994, which is a continuation-in-part of U.S. Serial No. 08/142,669 filed October 25, 1993, the contents of which are hereby incorporated by reference into the present disclosure.

Throughout this application, various publications are referred to by citations within parentheses and in the bibliographic description, immediately preceding the claims. The disclosures of these publications are hereby incorporated by reference into the present disclosure to more fully describe the state of the art to which this invention pertains.

Production of recombinant adenoviruses useful for gene therapy requires the use of a cell line capable of supplying in *trans* the gene products of the viral E1 region which are deleted in these recombinant viruses. At present the only useful cell line available is the 293 cell line originally described by Graham et al. in 1977. 293 cells contain approximately the left hand 12% (4.3 kb) of the adenovirus type 5 genome (Aiello (1979) and Spector (1983)).

Adenoviral vectors currently being tested for gene therapy applications typically are deleted for Ad2 or Ad5 DNA extending from approximately 400 base pairs from the 5' end of the viral genome to approximately 3.3 kb from the 5' end, for a total E1 deletion of 2.9 kb. Therefore, there exists a limited region of homology of approximately 1 kb between the DNA sequence of the recombinant virus and the Ad5 DNA within the cell line. This homology defines a region of potential recombination between the viral and cellular adenovirus sequences. Such a recombination results in a phenotypically wild-type virus bearing the Ad5 E1 region from the 293 cells. This recombination event

presumably accounts for the frequent detection of wild-type adenovirus in preparations of recombinant virus and has been directly demonstrated to be the cause of wild-type contamination of the Ad2 based recombinant virus Ad2/CFTR-1  
5 (Rich et al. (1993)).

Due to the high degree of sequence homology within the type C adenovirus subgroup such recombination is likely to occur if the vector is based on any group C adenovirus (types 1, 2, 5, 6).

10 In small scale production of recombinant adenoviruses, generation of contaminating wild-type virus can be managed by a screening process which discards those preparations of virus found to be contaminated. As the  
15 scale of virus production grows to meet expected demand for genetic therapeutics, the likelihood of any single lot being contaminated with a wild-type virus also will rise as well as the difficulty in providing non-contaminated recombinant preparations.

There will be over one million new cases of  
20 cancer diagnosed this year, and half that number of cancer-related deaths (American Cancer Society, 1993). p53 mutations are the most common genetic alteration associated with human cancers, occurring in 50-60% of human cancers (Hollstein et al. (1991); Bartek et al. (1991); Levine  
25 (1993)). The goal of gene therapy in treating p53 deficient tumors, for example, is to reinstate a normal, functional copy of the wild-type p53 gene so that control of cellular proliferation is restored. p53 plays a central role in cell cycle progression, arresting growth so that  
30 repair or apoptosis can occur in response to DNA damage. Wild-type p53 has recently been identified as a necessary component for apoptosis induced by irradiation or treatment with some chemotherapeutic agents (Lowe et al. (1993) A and B). Due to the high prevalence of p53 mutations in human

tumors, it is possible that tumors which have become refractory to chemotherapy and irradiation treatments may have become so due in part to the lack of wild-type p53. By resupplying functional p53 to these tumors, it is  
5 reasonable that they now are susceptible to apoptosis normally associated with the DNA damage induced by radiation and chemotherapy.

One of the critical points in successful human tumor suppressor gene therapy is the ability to affect a  
10 significant fraction of the cancer cells. The use of retroviral vectors has been largely explored for this purpose in a variety of tumor models. For example, for the treatment of hepatic malignancies, retroviral vectors have been employed with little success because these vectors are  
15 not able to achieve the high level of gene transfer required for *in vivo* gene therapy (Huber, B.E. et al., 1991; Caruso M. et al., 1993).

To achieve a more sustained source of virus production, researchers have attempted to overcome the  
20 problem associated with low level of gene transfer by direct injection of retroviral packaging cell lines into solid tumors (Caruso, M. et al., 1993; Ezzidine, Z.D. et al., 1991; Culver, K.W. et al., 1992). However, these methods are unsatisfactory for use in human patients  
25 because the method is troublesome and induces an inflammatory response against the packaging cell line in the patient. Another disadvantage of retroviral vectors is that they require dividing cells to efficiently integrate and express the recombinant gene of interest (Huber, B.E.  
30 1991). Stable integration into an essential host gene can lead to the development or inheritance of pathogenic diseased states.

Recombinant adenoviruses have distinct advantages over retroviral and other gene delivery methods (for

review, see Siegfried (1993)). Adenoviruses have never been shown to induce tumors in humans and have been safely used as live vaccines (Straus (1984)). Replication deficient recombinant adenoviruses can be produced by replacing the E1 region necessary for replication with the target gene. Adenovirus does not integrate into the human genome as a normal consequence of infection, thereby greatly reducing the risk of insertional mutagenesis possible with retrovirus or adeno-associated viral (AAV) vectors. This lack of stable integration also leads to an additional safety feature in that the transferred gene effect will be transient, as the extrachromosomal DNA will be gradually lost with continued division of normal cells. Stable, high titer recombinant adenovirus can be produced at levels not achievable with retrovirus or AAV, allowing enough material to be produced to treat a large patient population. Moreover, adenovirus vectors are capable of highly efficient *in vivo* gene transfer into a broad range of tissue and tumor cell types. For example, others have shown that adenovirus mediated gene delivery has a strong potential for gene therapy for diseases such as cystic fibrosis (Rosenfeld et al. (1992); Rich et al. (1993)) and  $\alpha_1$ -antitrypsin deficiency (Lemarchand et al. (1992)). Although other alternatives for gene delivery, such as cationic liposome/DNA complexes, are also currently being explored, none as yet appear as effective as adenovirus mediated gene delivery.

As with treating p53 deficient tumors, the goal of gene therapy for other tumors is to reinstate control of cellular proliferation. In the case of p53, introduction of a functional gene reinstates cell cycle control allowing for apoptotic cell death induced by therapeutic agents. Similarly, gene therapy is equally applicable to other tumor suppressor genes which can be used either alone or in combination with therapeutic agents to control cell cycle progression of tumor cells and/or induce cell death.

Moreover, genes which do not encode cell cycle regulatory proteins, but directly induce cell death such as suicide genes or, genes which are directly toxic to the cell can be used in gene therapy protocols to directly eliminate the cell cycle progression of tumor cells.

Regardless of which gene is used to reinstate the control of cell cycle progression, the rationale and practical applicability of this approach is identical. Namely, to achieve high efficiencies of gene transfer to express therapeutic quantities of the recombinant product. The choice of which vector to use to enable high efficiency gene transfer with minimal risk to the patient is therefore important to the level of success of the gene therapy treatment.

Thus, there exists a need for vectors and methods which provide high level gene transfer efficiencies and protein expression which provide safe and effective gene therapy treatments. The present invention satisfies this need and provides related advantages as well.

#### SUMMARY OF THE INVENTION

This invention provides a recombinant adenovirus expression vector characterized by the partial or total deletion of the adenoviral protein IX DNA and having a gene encoding a foreign protein or a functional fragment or mutant thereof. Transformed host cells and a method of producing recombinant proteins and gene therapy also are included within the scope of this invention.

Thus, for example, the adenoviral vector of this invention can contain a foreign gene for the expression of a protein effective in regulating the cell cycle, such as p53, Rb, or mitosin, or in inducing cell death, such as the conditional suicide gene thymidine kinase. (The latter

must be used in conjunction with a thymidine kinase metabolite in order to be effective).

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows a recombinant adenoviral vector of this invention. This construct was assembled as shown in Figure 1. The resultant virus bears a 5' deletion of adenoviral sequences extending from nucleotide 356 to 4020 and eliminates the E1a and E1b genes as well as the entire protein IX coding sequence, leaving the polyadenylation site shared by the E1b and pIX genes intact for use in terminating transcription of any desired gene.

Figure 2 shows the amino acid sequence of p110<sup>RB</sup>.

Figure 3 shows a DNA sequence encoding a retinoblastoma tumor suppressor protein.

Figure 4 shows schematic of recombinant p53/adenovirus constructs within the scope of this invention. The p53 recombinants are based on Ad 5 and have had the E1 region of nucleotides 360-3325 replaced with a 1.4 kb full length p53 cDNA driven by the Ad 2 MLP (A/M/53) or human CMV (A/C/53) promoters followed by the Ad 2 tripartite leader cDNA. The control virus A/M has the same Ad 5 deletions as the A/M/53 virus but lacks the 1.4 kb p53 cDNA insert. The remaining E1b sequence (705 nucleotides) have been deleted to create the protein IX deleted constructs A/M/N/53 and A/C/N/53. These constructs also have a 1.9 kb Xba I deletion within adenovirus type 5 region E3.

Figures 5A and 5B show p53 protein expression in tumor cells infected with A/M/53 and A/C/53. Figure 5A) Saos-2 (osteosarcoma) cells were infected at the indicated multiplicities of infection (MOI) with either the A/M/53 or



A/C/53 purified virus and harvested 24 hours later. The p53 antibody pAb 1801 was used to stain immunoblots of samples loaded at equal total protein concentrations. Equal protein concentration of SW480 cell extracts, which  
5 overexpress mutant p53 protein, were used as a marker for p53 size. "O" under the A/C/53 heading indicates a mock infection, containing untreated Saos-2 lysate. Figure 5B) Hep 3B (hepatocellular carcinoma) cells were infected with the A/M/53 or A/C/53 virus at the indicated MOI and  
10 analyzed as in part A.) The arrow indicates the position of the p53 protein.

Figures 6A through 6C show p53 dependent Saos-2 morphology change. Subconfluent ( $1 \times 10^5$  cells/10 cm plate) Saos-2 cells were either uninfected (A), infected at an MOI  
15 = 50 with (B) the control A/M virus or (C) the A/C/53 virus. The cells were photographed 72 hours post-infection.

Figure 7 shows p53 dependent inhibition of DNA synthesis in human tumor cell lines by A/M/N/53 and  
20 A/C/N/53. Nine different tumor cell lines were infected with either control adenovirus A/M (-x-x-), or the p53 expressing A/M/N/53 (-Δ-Δ-), or A/C/N/53 (-O-O-) virus at increasing MOI as indicated. The tumor type and p53 status is noted for each cell line (wt = wild type, null = no  
25 protein expressed, mut = mutant protein expressed). DNA synthesis was measured 72 hours post-infection as described below in Experiment No. II. Results are from triplicate measurements at each dose (mean+/- SD), and are plotted as % of media control versus MOI. \* H69 cells were only  
30 tested with A/M and A/M/N/53 virus.

Figure 8 shows tumorigenicity of p53 infected Saos-2 cells in nude mice. Saos-2 cells were infected with either the control A/M virus or the p53 recombinant A/M/N/53 at MOI = 30. Treated cells were injected

2025 RELEASE UNDER E.O. 14176

subcutaneously into the flanks of nude mice, and tumor dimensions were measured (as described in Experiment No. II) twice per week for 8 weeks. Results are plotted as tumor size versus days post tumor cell implantation for both control A/M (-x-x-) and A/M/N/53 (-Δ-Δ-) treated cells. Error bars represent the mean tumor size  $\pm$  SEM for each group of 4 animals at each time point.

Figure 9 is expression of rAd/p53 RNA in established tumors. H69 (SCLC) cells were injected subcutaneously into nude mice and allowed to develop tumors for 32 days until reaching a size of approximately 25-50 mm<sup>3</sup>. Mice were randomized and injected peritumorally with  $2 \times 10^9$  pfu of either control A/C/β-gal or A/C/53 virus. Tumors were excised 2 and 7 days post injection, and polyA RNA was prepared from each tumor sample. RT-PCR was carried out using equal RNA concentrations and primers specific for recombinant p53 message. PCR amplification was for 30 cycles at 94°C 1 min., 55°C 1.5 min., 72°C 2 min., and a 10 min., 72°C final extension period in an Omnigen thermalcycler (Hybaid). The PCR primers used were a 5' Tripartite Leader cDNA (5' - CGCCACCGAGGGACCTGAGCGAGTC-3') and a 3' p53 primer (5' - TTCTGGGAAGGGACAGAAGA-3'). Lanes 1, 2, 4, and 5 are p53 treated samples excised at day 2 or 7 as indicated. Lanes 3 and 6 are from β-gal treated tumors. Lanes 7,8, and 9 are replicates of lanes 4,5, and 6 respectively, amplified with actin primers to verify equal loading. Lane 10 is a positive control using a tripartite/p53 containing plasmid.

Figures 10A and 10B show *in vivo* tumor suppression and increased survival time with A/M/N/53. H69 (SCLC) tumor cells were injected subcutaneously into nude mice and allowed to develop for 2 weeks. Peritumoral injections of either buffer alone (---), control A/M adenovirus (-x-x-), or A/M/N/53 (-Δ-Δ), both viruses ( $2 \times 10^9$  pfu/injection) were administered twice per week for a

total of 8 doses. Tumor dimensions were measured twice per week and tumor volume was estimated as described in Experiment No. II. A) Tumor size is plotted for each virus versus time (days) post inoculation of H69 cells. Error bars indicate the mean tumor size +/- SEM for each group of 5 animals. Arrows indicate days virus injections. B) Mice were monitored for survival and the fraction of mice surviving per group versus time post inoculation of buffer alone (----), control A/M (... ..) or A/M/N/53 (—) virus treated H69 cells is plotted.

Figures 11A through 11C show maps of recombinant plasmid constructions. Plasmids were constructed as detailed in below. Bold lines in the constructs indicate genes of interest while boldface type indicates the restriction sites used to generate the fragments to be ligated together to form the subsequent plasmid as indicated by the arrows. In Figure 11A, the plasmid pACNTK was constructed by subcloning the HSV-TK gene from pMLBTK (ATCC No. 39369) into the polylinker of a cloning vector, followed by isolation of the TK gene with the desired ends for cloning into the pACN vector. The pACN vector contains adenoviral sequences necessary for *in vivo* recombination to occur to form recombinant adenovirus (see Figure 12). In Figure 11B, the construction of the plasmid pAANTK is shown beginning with PCR amplified fragments encoding the  $\alpha$ -fetoprotein enhancer (AFP-E) and promoter (AFP-P) regions subcloned through several steps into a final plasmid where the AFP enhancer and promoter are upstream of the HSV-TK gene followed by adenovirus Type 2 sequences necessary for *in vivo* recombination to occur to form recombinant adenovirus. In Figure 11C, the construction of the plasmid pAANCAT is shown beginning with the isolation of the chloramphenicol acetyltransferase (CAT) gene from a commercially available plasmid and subcloning it into the pAAN plasmid (see above), generating the final plasmid pAANCAT where the AFP enhancer/promoter direct

transcription of the CAT gene in an adenovirus sequence background.

Figure 12 is a schematic map of recombinant adenoviruses ACNTK, AANTK and AANCAT. To construct  
5 recombinant adenoviruses from the plasmids described in Figure 11, 4 parts (20  $\mu$ g) of either plasmid pACNTK, pAANTK, or pAANCAT were linearized with Eco R1 and cotransfected with 1 part (5  $\mu$ g) of the large fragment of  
10 Cla 1 digested recombinant adenovirus (rAC $\beta$ -gal) containing an E3 region deletion (Wills et al., 1994). In the resulting viruses, the Ad 5 nucleotides 360 - 4021 are replaced by either the CMV promoter and tripartite leader cDNA (TPL) or the  $\alpha$ -fetoprotein enhancer and promoter (AFP) driving expression of the HSV-1 TK or CAT gene as  
15 indicated. The resulting recombinant adenoviruses are designated ACNTK, AANTK, and AANCAT respectively.

Figure 13 shows promoter specificity of CAT expression in the recombinant adenoviral vectors. Two (2)  $\times 10^6$  of the designated cell lines were infected at MOIs =  
20 30 or 100 of the recombinant adenovirus AANCAT as indicated or left uninfected (UN). Hep G2 and Hep 3B cells express  $\alpha$ -fetoprotein whereas the other cell lines do not. After three days, the cells were harvested, extract volumes were adjusted for equal total protein concentrations, and CAT  
25 activity was measured as described in Methods section, below. An equal number of uninfected cells served as individual controls for background CAT activity, while  $^{14}$ C labelled chloramphenicol ( $^{14}$ C-only) and extract from a stable cell line (B21) expressing CAT activity served as  
30 negative and positive controls respectively. Percent conversion of acetyl CoA is indicated, demonstrating that CAT expression is limited to those cells expressing  $\alpha$ -fetoprotein.

Figure 14 shows the effects of TK/GCV treatment on hepatocellular carcinoma cell lines and the effects of promoter specificity. Hep-G2 (AFP positive) and HLF (AFP negative) cell lines were infected overnight with ACNTK [- $\Delta$ -] AANTK [- $\Delta$ -], or control ACN [- $\square$ -] virus at an infection multiplicity of 30 and subsequently treated with a single dose of ganciclovir at the indicated concentrations. Cell proliferation was assessed by adding  $^3\text{H}$ -thymidine to the cells approximately 18 hours prior to harvest.  $^3\text{H}$ -thymidine incorporation into cellular nucleic acid was measured 72 hours after infection (Top Count, Packard and expressed as a percent (mean  $\pm$  S.D.) of untreated control. The results show a non-selective dose dependent inhibition of proliferation with the CMV driven construct, while AFP driven TK selectively inhibits Hep-G2.

Figure 15 shows cytotoxicity of ACNTK plus ganciclovir in HCC. HLF cells were infected at an MOI of 30 with either ACNTK [- $\bullet$ -] or the control virus ACN [- $\square$ -] and treated with ganciclovir at the indicated doses. Seventy-two (72) hours after ganciclovir treatment, the amount of lactate dehydrogenase (LDH) released into the cell supernatant were measured colorimetrically and plotted (mean $\pm$ -SEM) versus ganciclovir concentration for the two virus treated groups.

Figures 16A and 16B show the effect of ACNTK plus ganciclovir on established hepatocellular carcinoma (HCC) tumors in nude mice. One ( $1 \times 10^7$ ) Hep 3B cells were injected subcutaneously into the flank of female nude mice and allowed to grow for 27 days. Mice then received intratumoral and peritumoral injections of either the ACNTK [- $\bullet$ -] or control ACN [- $\square$ -] virus ( $1 \times 10^9$  iu in 100  $\mu\text{l}$  volume) every other day for a total of three doses (indicated by arrows). Injections of ganciclovir (100 mg/kg ip) began 24 hours after the initial virus dose and continued for a total of 10 days. In Figure 6A, tumor

sizes are plotted for each virus versus days post infection (mean +/- SEM). In Figure 6B, body weight for each virus-treated animal group is plotted as the mean +/-SEM versus days post infection.

5

#### DETAILED DESCRIPTION OF THE INVENTION

To reduce the frequency of contamination with wild-type adenovirus, it is desirable to improve either the virus or the cell line to reduce the probability of recombination. For example, an adenovirus from a group  
10 with low homology to the group C viruses could be used to engineer recombinant viruses with little propensity for recombination with the Ad5 sequences in 293 cells. However, an alternative, easier means of reducing the recombination between viral and cellular sequences is to  
15 increase the size of the deletion in the recombinant virus and thereby reduce the extent of shared sequence between it and the Ad5 genes in the 293 cells.

Deletions which extend past 3.5 kb from the 5' end of the adenoviral genome affect the gene for adenoviral  
20 protein IX and have not been considered desirable in adenoviral vectors (see below).

The protein IX gene of the adenoviruses encodes a minor component of the outer adenoviral capsid which stabilizes the group-of-nine hexons which compose the  
25 majority of the viral capsid (Stewart (1993)). Based upon study of adenovirus deletion mutants, protein IX initially was thought to be a non-essential component of the adenovirus, although its absence was associated with greater heat lability than observed with wild-type virus  
30 (Colby and Shenk (1981)). More recently it was discovered that protein IX is essential for packaging full length viral DNA into capsids and that in the absence of protein IX, only genomes at least 1 kb smaller than wild-type could

be propagated as recombinant viruses (Ghosh-Choudhury et al. (1987)). Given this packaging limitation, protein IX deletions deliberately have not been considered in the design of adenoviral vectors.

5           In this application, reference is made to standard textbooks of molecular biology that contain definitions, methods and means for carrying out basic techniques, encompassed by the present invention. See for example, Sambrook et al. (1989) and the various references  
10   cited therein. This reference and the cited publications are expressly incorporated by reference into this disclosure.

Contrary to what has been known in the art, this invention claims the use of recombinant adenoviruses  
15   bearing deletions of the protein IX gene as a means of reducing the risk of wild-type adenovirus contamination in virus preparations for use in diagnostic and therapeutic applications such as gene therapy. As used herein, the term "recombinant" is intended to mean a progeny formed as  
20   the result of genetic engineering. These deletions can remove an additional 500 to 700 base pairs of DNA sequence that is present in conventional E1 deleted viruses (smaller, less desirable, deletions of portions of the pIX gene are possible and are included within the scope of this  
25   invention) and is available for recombination with the Ad5 sequences integrated in 293 cells. Recombinant adenoviruses based on any group C virus, serotype 1, 2, 5 and 6, are included in this invention. Also encompassed by this invention is a hybrid Ad2/Ad5 based recombinant virus  
30   expressing the human p53 cDNA from the adenovirus type 2 major late promoter. This construct was assembled as shown in Figure 1. The resultant virus bears a 5' deletion of adenoviral sequences extending from about nucleotide 357 to 4020 and eliminates the E1a and E1b genes as well as the  
35   entire protein IX coding sequence, leaving the

polyadenylation site shared by the Elb and protein IX genes intact for use in terminating transcription of any desired gene. A separate embodiment is shown in Figure 4. Alternatively, the deletion can be extended an additional 5 30 to 40 base pairs without affecting the adjacent gene for protein IVa2, although in that case an exogenous polyadenylation signal is provided to terminate transcription of genes inserted into the recombinant virus. The initial virus constructed with this deletion is easily 10 propagated in 293 cells with no evidence of wild-type viral contamination and directs robust p53 expression from the transcriptional unit inserted at the site of the deletion.

The insert capacity of recombinant viruses bearing the protein IX deletion described above is 15 approximately 2.6 kb. This is sufficient for many genes including the p53 cDNA. Insert capacity can be increased by introducing other deletions into the adenoviral backbone, for example, deletions within early regions 3 or 4 (for review see: Graham and Prevec (1991)). For 20 example, the use of an adenoviral backbone containing a 1.9 kb deletion of non-essential sequence within early region 3. With this additional deletion, the insert capacity of the vector is increased to approximately 4.5 kb, large enough for many larger cDNAs, including that of the 25 retinoblastoma tumor suppressor gene.

A recombinant adenovirus expression vector characterized by the partial or total deletion of the adenoviral protein IX DNA and having a gene encoding a foreign protein, or a functional fragment or mutant thereof 30 is provided by this invention. These vectors are useful for the safe recombinant production of diagnostic and therapeutic polypeptides and proteins, and more importantly, for the introduction of genes in gene therapy. Thus, for example, the adenoviral vector of this invention 35 can contain a foreign gene for the expression of a protein



effective in regulating the cell cycle, such as p53, Rb, or mitotin, or in inducing cell death, such as the conditional suicide gene thymidine kinase. (The latter must be used in conjunction with a thymidine kinase metabolite in order to be effective). Any expression cassette can be used in the vectors of this invention. An "expression cassette" means a DNA molecule having a transcription promoter/enhancer such as the CMV promoter enhancer, etc., a foreign gene, and in some embodiments defined below, a polyadenylation signal. As used herein, the term "foreign gene" is intended to mean a DNA molecule not present in the exact orientation and position as the counterpart DNA molecule found in wild-type adenovirus. The foreign gene is a DNA molecule up to 4.5 kilobases. "Expression vector" means a vector that results in the expression of inserted DNA sequences when propagated in a suitable host cell, i.e., the protein or polypeptide coded for by the DNA is synthesized by the host's system. The recombinant adenovirus expression vector can contain part of the gene encoding adenovirus protein IX, provided that biologically active protein IX or fragment thereof is not produced. Example of this vector are an expression vector having the restriction enzyme map of Figures 1 or 4.

Inducible promoters also can be used in the adenoviral vector of this invention. These promoters will initiate transcription only in the presence of an additional molecule. Examples of inducible promoters include those obtainable from a  $\beta$ -interferon gene, a heat shock gene, a metallothioneine gene or those obtainable from steroid hormone-responsive genes. Tissue specific expression has been well characterized in the field of gene expression and tissue specific and inducible promoters such as these are very well known in the art. These genes are used to regulate the expression of the foreign gene after it has been introduced into the target cell.

Also provided by this invention is a recombinant adenovirus expression vector, as described above, having less extensive deletions of the protein IX gene sequence extending from 3500 bp from the 5' viral termini to approximately 4000 bp, in one embodiment. In a separate embodiment, the recombinant adenovirus expression vector can have a further deletion of a non-essential DNA sequence in adenovirus early region 3 and/or 4 and/or deletion of the DNA sequences designated adenovirus E1a and E1b. In this embodiment, foreign gene is a DNA molecule of a size up to 4.5 kilobases.

A further embodiment has a deletion of up to forty nucleotides positioned 3' to the E1a and E1b deletion and pIX and a foreign DNA molecule encoding a polyadenylation signal inserted into the recombinant vector in a position relative to the foreign gene to regulate the expression of the foreign gene.

For the purposes of this invention, the recombinant adenovirus expression vector can be derived from wild-type group adenovirus, serotype 1, 2, 5 or 6.

In one embodiment, the recombinant adenovirus expression vector has a foreign gene coding for a functional tumor suppressor protein, or a biologically active fragment thereof. As used herein, the term "functional" as it relates to a tumor suppressor gene, refers to tumor suppressor genes that encode tumor suppressor proteins that effectively inhibit a cell from behaving as a tumor cell. Functional genes can include, for instance, wild type of normal genes and modifications of normal genes that retains its ability to encode effective tumor suppressor proteins and other anti-tumor genes such as a conditional suicide protein or a toxin.

Similarly, "non-functional" as used herein is synonymous with "inactivated." Non-functional or defective genes can be caused by a variety of events, including for example point mutations, deletions, methylation and others known to those skilled in the art.

As used herein, an "active fragment" of a gene includes smaller portions of the gene that retain the ability to encode proteins having tumor suppressing activity. p56<sup>RB</sup>, described more fully below, is but one example of an active fragment of a functional tumor suppressor gene. Modifications of tumor suppressor genes are also contemplated within the meaning of an active fragment, such as additions, deletions or substitutions, as long as the functional activity of the unmodified gene is retained.

Another example of a tumor suppressor gene is retinoblastoma (RB). The complete RB cDNA nucleotide sequences and predicted amino acid sequences of the resulting RB protein (designated p110<sup>RB</sup>) are shown in Lee et al. (1987) and in Figure 3. Also useful to express retinoblastoma tumor suppressor protein is a DNA molecule encoding the amino acid sequence shown in Figure 2 or having the DNA sequence shown in Figure 3. A truncated version of p110<sup>RB</sup>, called p56<sup>RB</sup> also is useful. For the sequence of p56<sup>RB</sup>, see Huang et al. (1991). Additional tumor suppressor genes can be used in the vectors of this invention. For illustration purposes only, these can be p16 protein (Kamb et al. (1994)), p21 protein, Wilm's tumor WT1 protein, mitosin, h-NUC, or colon carcinoma DCC protein. Mitosin is described in X. Zhu and W-H Lee, U.S. Application Serial No. 08/141,239, filed October 22, 1993, and a subsequent continuation-in-part by the same inventors, attorney docket number P-CJ 1191, filed October 24, 1994, both of which are herein incorporated by reference. Similarly, h-NUC is described by W-H Lee and P-

L Chen, U.S. Application Serial No. 08/170,586, filed December 20, 1993, herein incorporated by reference.

As is known to those of skill in the art, the term "protein" means a linear polymer of amino acids joined in a specific sequence by peptide bonds. As used herein, the term "amino acid" refers to either the D or L stereoisomer form of the amino acid, unless otherwise specifically designated. Also encompassed within the scope of this invention are equivalent proteins or equivalent peptides, e.g., having the biological activity of purified wild type tumor suppressor protein. "Equivalent proteins" and "equivalent polypeptides" refer to compounds that depart from the linear sequence of the naturally occurring proteins or polypeptides, but which have amino acid substitutions that do not change its biologically activity. These equivalents can differ from the native sequences by the replacement of one or more amino acids with related amino acids, for example, similarly charged amino acids, or the substitution or modification of side chains or functional groups.

Also encompassed within the definition of a functional tumor suppressor protein is any protein whose presence reduces the tumorigenicity, malignancy or hyperproliferative phenotype of the host cell. Examples of tumor suppressor proteins within this definition include, but are not limited to p110<sup>RB</sup>, p56<sup>RB</sup>, mitosin, h-NUC and p53. "Tumorigenicity" is intended to mean having the ability to form tumors or capable of causing tumor formation and is synonymous with neoplastic growth. "Malignancy" is intended to describe a tumorigenic cell having the ability to metastasize and endanger the life of the host organism. "Hyperproliferative phenotype" is intended to describe a cell growing and dividing at a rate beyond the normal limitations of growth for that cell type. "Neoplastic" also is intended to include cells lacking endogenous

functional tumor suppressor protein or the inability of the cell to express endogenous nucleic acid encoding a functional tumor suppressor protein.

5 An example of a vector of this invention is a recombinant adenovirus expression vector having a foreign gene coding for p53 protein or an active fragment thereof is provided by this invention. The coding sequence of the p53 gene is set forth below in Table I.

2595370-102597

TABLE 1

50

V\*SHR PGSR\* LLGSG DTLRS GWER A FHDGD TLPWI GSQTA FRVTA MEEPQ  
 100

5 SDPSV EPPLS QETFS DLWKL LPENN VLSPL PSQAM DDLML SPDDI EQWFT  
 150

EDPGP DEAPR MPEAA PPVAP APAAP TPAAP APAPS WPLSS SVPSQ KTYQG  
 200

SYGFR LGFLH SGTAK SVTCT YSPAL NKMFC QLAKT CPVQL WVDST PPPGT  
 10 250

RVRAM AIYKQ SQHMT EVVRR CPHHE RCSDS DGLAP PQHLI RVEGN LRVEY  
 300

LDDRN TFRHS VVVPY EPPEV GSDCT TIHYN YMCNS SCMGG MNRRP ILTII  
 350

15 TLED SGNLL GRNSF EVRVC ACPGR DRRTE EENLR KKGEP HHELP PGSTK  
 400

RALPN NTSSS PQPKK KPLDG EYFTL QIRGR ERFEM FRELN EALEL KDAQA

GKEPG GSRAH SSHLK SKKGQ STSRH KKLMF KTEGP DSD\*

\* = Stop codon

20 Any of the expression vectors described herein  
 are useful as compositions for diagnosis or therapy. The  
 vectors can be used for screening which of many tumor  
 suppressor genes would be useful in gene therapy. For  
 example, a sample of cells suspected of being neoplastic  
 25 can be removed from a subject and mammal. The cells can  
 then be contacted, under suitable conditions and with an  
 effective amount of a recombinant vector of this invention  
 having inserted therein a foreign gene encoding one of  
 several functional tumor suppressor genes. Whether the  
 30 introduction of this gene will reverse the malignant  
 phenotype can be measured by colony formation in soft agar  
 or tumor formation in nude mice. If the malignant  
 phenotype is reversed, then that foreign gene is determined  
 to be a positive candidate for successful gene therapy for  
 35 the subject or mammal. When used pharmaceutically, they

5

15

20

25

30

35

pharmaceutical composition also can be incorporated, if desired, into liposomes, microspheres or other polymer matrices (Gregoriadis, Liposome Technology, Vol. 1 (CRC Press, Boca Raton, Florida 1984), which is incorporated  
5 herein by reference). Liposomes, for example, which consist of phospholipids or other lipids, are nontoxic, physiologically acceptable and metabolizable carriers that are relatively simple to make and administer.

As used herein, "pharmaceutical composition"  
10 refers to any of the compositions of matter described herein in combination with one or more of the above pharmaceutically acceptable carriers. The compositions can then be administered therapeutically or prophylactically. They can be contacted with the host cell *in vivo*, *ex vivo*,  
15 or *in vitro*, in an effective amount. *In vitro* and *ex vivo* means of contacting host cells are provided below. When practiced *in vivo*, methods of administering a pharmaceutical containing the vector of this invention, are well known in the art and include but are not limited to,  
20 administration orally, intra-tumorally, intravenously, intramuscularly or intraperitoneal. Administration can be effected continuously or intermittently and will vary with the subject and the condition to be treated, e.g., as is the case with other therapeutic compositions (Landmann et  
25 al. (1992); Aulitzky et al. (1991); Lantz et al. (1990); Supersaxo et al. (1988); Demetri et al. (1989); and LeMaistre et al. (1991)).

Further provided by this invention is a transformed procaryotic or eucaryotic host cell, for  
30 example an animal cell or mammalian cell, having inserted a recombinant adenovirus expression vector described above. Suitable procaryotic cells include but are not limited to bacterial cells such as *E. coli* cells. Methods of transforming host cells with retroviral vectors are known  
35 in the art, see Sambrook et al. (1989) and include, but are



not limited to transfection, electroporation, and microinjection.

As used throughout this application, the term animal is intended to be synonymous with mammal and is to include, but not be limited to bovine, porcine, feline, simian, canine, equine, murine, rat or human. Additional host cells include but are not limited to any neoplastic or tumor cell, such as osteosarcoma, ovarian carcinoma, breast carcinoma, melanoma, hepatocarcinoma, lung cancer, brain cancer, colorectal cancer, hematopoietic cell, prostate cancer, cervical carcinoma, retinoblastoma, esophageal carcinoma, bladder cancer, neuroblastoma, or renal cancer.

Additionally, any eucaryotic cell line capable of expressing Ela and Elb or Ela, Elb and pIX is a suitable host for this vector. In one embodiment, a suitable eucaryotic host cell is the 293 cell line available from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland, U.S.A. 20231.

Any of the transformed host cells described herein are useful as compositions for diagnosis or therapy. When used pharmaceutically, they can be combined with various pharmaceutically acceptable carriers. Suitable pharmaceutically acceptable carriers are well known to those of skill in the art and, for example, are described above. The compositions can then be administered therapeutically or prophylactically, in effective amounts, described in more detail below.

A method of transforming a host cell also is provided by this invention. This method provides contacting a host cell, i.e., a procaryotic or eucaryotic host cell, with any of the expression vectors described herein and under suitable conditions. Host cells transformed by this method also are claimed within the

scope of this invention. The contacting can be effected *in vitro*, *in vivo*, or *ex vivo*, using methods well known in the art (Sambrook et al. (1989)) and using effective amounts of the expression vectors. Also provided in this invention is

5 a method of producing a recombinant protein or polypeptide by growing the transformed host cell under suitable conditions favoring the transcription and translation of the inserted foreign gene. Methods of recombinant expression in a variety of host cells, such as mammalian,

10 yeast, insect or bacterial cells, are widely known, including those described in Sambrook et al., supra. The translated foreign gene can then be isolated by convention means, such as column purification or purification using an anti-protein antibody. The isolated protein or polypeptide

15 also is intended within the scope of this invention. As used herein, purified or isolated mean substantially free of native proteins or nucleic acids normally associated with the protein or polypeptide in the native or host cell environment.

20 Also provided by this invention are non-human animals having inserted therein the expression vectors or transformed host cells of this invention. These "transgenic" animals are made using methods well known to those of skill in the art, for example as described in U.S.

25 Patent No. 5,175,384 or by conventional *ex vivo* therapy techniques, as described in Culver et al. (1991).

As shown in detail below, the recombinant adenoviruses expressing a tumor suppressor wild-type p53, as described above, can efficiently inhibit DNA synthesis

30 and suppress the growth of a broad range of human tumor cell types, including clinical targets. Furthermore, recombinant adenoviruses can express tumor suppression genes such as p53 in an *in vivo* established tumor without relying on direct injection into the tumor or prior *ex vivo*

35 treatment of the cancer cells. The p53 expressed is

functional and effectively suppresses tumor growth *in vivo* and significantly increases survival time in a nude mouse model of human lung cancer.

Thus, the vectors of this invention are particularly suited for gene therapy. Accordingly, methods of gene therapy utilizing these vectors are within the scope of this invention. The vector is purified and then an effective amount is administered *in vivo* or *ex vivo* into the subject. Methods of gene therapy are well known in the art, see, for example, Larrick, J.W. and Burck, K.L. (1991) and Kreigler, M. (1990). "Subject" means any animal, mammal, rat, murine, bovine, porcine, equine, canine, feline or human patient. When the foreign gene codes for a tumor suppressor gene or other anti-tumor protein, the vector is useful to treat or reduce hyperproliferative cells in a subject, to inhibit tumor proliferation in a subject or to ameliorate a particular related pathology. Pathologic hyperproliferative cells are characteristic of the following disease states, thyroid hyperplasia - Grave's Disease, psoriasis, benign prostatic hypertrophy, Li-Fraumeni syndrome including breast cancer, sarcomas and other neoplasms, bladder cancer, colon cancer, lung cancer, various leukemias and lymphomas. Examples of non-pathologic hyperproliferative cells are found, for instance, in mammary ductal epithelial cells during development of lactation and also in cells associated with wound repair. Pathologic hyperproliferative cells characteristically exhibit loss of contact inhibition and a decline in their ability to selectively adhere which implies a change in the surface properties of the cell and a further breakdown in intercellular communication. These changes include stimulation to divide and the ability to secrete proteolytic enzymes.

Moreover, the present invention relates to a method for depleting a suitable sample of pathologic

1 mammalian hyperproliferative cells contaminating  
hematopoietic precursors during bone marrow reconstitution  
via the introduction of a wild type tumor suppressor gene  
into the cell preparation using the vector of this  
5 invention (whether derived from autologous peripheral blood  
or bone marrow). As used herein, a "suitable sample" is  
defined as a heterogeneous cell preparation obtained from  
a patient, e.g., a mixed population of cells containing  
both phenotypically normal and pathogenic cells.  
10 "Administer" includes, but is not limited to introducing  
into the cell or subject intravenously, by direct injection  
into the tumor, by intra-tumoral injection, by  
intraperitoneal administration, by aerosol administration  
to the lung or topically. Such administration can be  
15 combined with a pharmaceutically-accepted carrier,  
described above.

2 The term "reduced tumorigenicity" is intended to  
mean tumor cells that have been converted into less  
tumorigenic or non-tumorigenic cells. Cells with reduced  
20 tumorigenicity either form no tumors *in vivo* or have an  
extended lag time of weeks to months before the appearance  
of *in vivo* tumor growth and/or slower growing three  
dimensional tumor mass compared to tumors having fully  
inactivated or non-functional tumor suppressor gene.

25 As used herein, the term "effective amount" is  
intended to mean the amount of vector or anti-cancer  
protein which achieves a positive outcome on controlling  
cell proliferation. For example, one dose contains from  
about  $10^8$  to about  $10^{13}$  infectious units. A typical course  
30 of treatment would be one such dose a day over a period of  
five days. An effective amount will vary on the pathology  
or condition to be treated, by the patient and his status,  
and other factors well known to those of skill in the art.  
Effective amounts are easily determined by those of skill  
35 in the art.

This invention also provides a method for reducing the proliferation of tumor cells in a subject by introducing into the tumor mass an effective amount of an adenoviral expression vector containing an anti-tumor gene other than a tumor suppressor gene. The anti-tumor gene can encode, for example, thymidine kinase (TK). The subject is then administered an effective amount of a therapeutic agent, which in the presence of the anti-tumor gene is toxic to the cell. In the specific case of thymidine kinase, the therapeutic agent is a thymidine kinase metabolite such as ganciclovir (GCV), 6-methoxypurine arabinonucleoside (araM), or a functional equivalent thereof. Both the thymidine kinase gene and the thymidine kinase metabolite must be used concurrently to be toxic to the host cell. However, in its presence, GCV is phosphorylated and becomes a potent inhibitor of DNA synthesis whereas araM gets converted to the cytotoxic anabolite araATP. Other anti-tumor genes can be used as well in combination with the corresponding therapeutic agent to reduce the proliferation of tumor cells. Such other gene and therapeutic agent combinations are known by one skilled in the art. Another example would be the vector of this invention expressing the enzyme cytosine deaminase. Such vector would be used in conjunction with administration of the drug 5-fluorouracil (Austin and Huber, 1993), or the recently described E. Coli Deo  $\Delta$  gene

in combination with 6-methyl-purine-2'-deosribonucleoside (Sorscher et al 1994).

As with the use of the tumor suppressor genes described previously, the use of other anti-tumor genes, either alone or in combination with the appropriate therapeutic agent provides a treatment for the uncontrolled cell growth or proliferation characteristic of tumors and malignancies. Thus, this invention provides a therapy to stop the uncontrolled cellular growth in the patient thereby alleviating the symptoms of the disease or cachexia present in the patient. The effect of this treatment includes, but is not limited to, prolonged survival time of the patient, reduction in tumor mass or burden, apoptosis of tumor cells or the reduction of the number of circulating tumor cells. Means of quantifying the beneficial effects of this therapy are well known to those of skill in the art.

The invention provides a recombinant adenovirus expression vector characterized by the partial or total deletion of the adenoviral protein IX DNA and having a foreign gene encoding a foreign protein, wherein the foreign protein is a suicide gene or functional equivalent thereof. The anti-cancer gene TK, described above, is an example of a suicide gene because when expressed, the gene product is, or can be made to be lethal to the cell. For TK, lethality is induced in the presence of GCV. The TK gene is derived from herpes simplex virus by methods well known to those of skill in the art. The plasmid pMLBKTK in E. coli HB101 (from ATCC #39369) is a source of the herpes simplex virus (HSV-1) thymidine kinase (TK) gene for use in this invention. However, many other sources exist as well.

The TK gene can be introduced into the tumor mass by combining the adenoviral expression vector with a

00953570-102897  
268207 0458580

suitable pharmaceutically acceptable carrier. Introduction can be accomplished by, for example, direct injection of the recombinant adenovirus into the tumor mass. For the specific case of a cancer such as hepatocellular carcinoma (HCC), direct injection into the hepatic artery can be used for delivery because most HCCs derive their circulation from this artery. To control proliferation of the tumor, cell death is induced by treating the patients with a TK metabolite such as ganciclovir to achieve reduction of tumor mass. The TK metabolite can be administered, for example, systemically, by local innoculation into the tumor or in the specific case of HCC, by injection into the hepatic artery. The TK metabolite is preferably administered at least once daily but can be increased or decreased according to the need. The TK metabolite can be administered simultaneous or subsequent to the administration of the TK containing vector. Those skilled in the art know or can determine the dose and duration which is therapeutically effective.

20           A method of tumor-specific delivery of a tumor suppressor gene is accomplished by contacting target tissue in an animal with an effective amount of the recombinant adenoviral expression vector of this invention. The gene is intended to code for an anti-tumor agent, such as a functional tumor suppressor gene or suicide gene. "Contacting" is intended to encompass any delivery method for the efficient transfer of the vector, such as intra-tumoral injection.

30           The use of the adenoviral vector of this invention to prepare medicaments for the treatment of a disease or for therapy is further provided by this invention.

          The following examples are intended to illustrate, not limit the scope of this invention.

## EXPERIMENT NO. I

Plasmid pAd/MLP/p53/E1b- was used as the starting material for these manipulations. This plasmid is based on the pBR322 derivative pML2 (pBR322 deleted for base pairs 1140 to 2490) and contains adenovirus type 5 sequences extending from base pair 1 to base pair 5788 except that it is deleted for adenovirus type 5 base pairs 357 to 3327. At the site of the Ad5 357/3327 deletion a transcriptional unit is inserted which is comprised of the adenovirus type 2 major late promoter, the adenovirus type 2 tripartite leader cDNA and the human p53 cDNA. It is a typical E1 replacement vector deleted for the Ad5 E1a and E1b genes but containing the Ad5 protein IX gene (for review of Adenovirus vectors see: Graham and Prevec (1992)). Ad2 DNA was obtained from Gibco BRL. Restriction endonucleases and T4 DNA ligase were obtained from New England Biolabs. *E. coli* DH5 $\alpha$  competent cells were purchased from Gibco BRL and 293 cells were obtained from the American Type Culture Collection (ATCC). Prep-A-Gene DNA purification resin was obtained from BioRad. LB broth bacterial growth medium was obtained from Difco. Qiagen DNA purification columns were obtained from Qiagen, Inc. Ad5 dl327 was obtained from R.J. Schneider, NYU. The MBS DNA transfection kit was purchased from Stratagene.

One (1)  $\mu$ g pAd/MLP/p53/E1b- was digested with 20 units each of restriction enzymes *Ecl* 136II and *Ngo*MI according to the manufacturer's recommendations. Five (5)  $\mu$ g Ad2 DNA was digested with 20 units each of restriction endonucleases *Dra*I and *Ngo*MI according to the manufacturer's recommendations. The restriction digestions were loaded into separate lanes of a 0.8% agarose gel and electrophoresed at 100 volts for 2 hours. The 4268 bp restriction fragment from the Pad/MLP/p53/E1b- sample and the 6437 bp fragment from the Ad2 sample were isolated from the gel using Prep-A-Gene DNA extraction resin according to



the manufacturer's specifications. The restriction fragments were mixed and treated with T4 DNA ligase in a total volume of 50  $\mu$ l at 16°C for 16 hours according to the manufacturer's recommendations. Following ligation 5  $\mu$ l of the reaction was used to transform *E. coli* DH5 $\alpha$  cells to ampicillin resistance following the manufacturer's procedure. Six bacterial colonies resulting from this procedure were used to inoculate separate 2 ml cultures of LB growth medium and incubated overnight at 37°C with shaking. DNA was prepared from each bacterial culture using standard procedures (Sambrook et al. (1989)). One fourth of the plasmid DNA from each isolate was digested with 20 units of restriction endonuclease XhoI to screen for the correct recombinant containing XhoI restriction fragments of 3627, 3167, 2466 and 1445 base pairs. Five of six screened isolates contained the correct plasmid. One of these was then used to inoculate a 1 liter culture of LB medium for isolation of large quantities of plasmid DNA. Following overnight incubation plasmid DNA was isolated from the 1 liter culture using Qiagen DNA purification columns according to the manufacturer's recommendations. The resulting plasmid was designated Pad/MLP/p53/PIX-. Samples of this plasmid were deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland, U.S.A., 12301, on October 22, 1993. The deposit was made under the provisions of the Budapest Treaty on the International Deposit of Microorganisms for the Purpose of Patent Procedure. The deposit was accorded ATCC Accession No. 75576.

To construct a recombinant adenovirus, 10  $\mu$ g Pad/MLP/p53/PIX- were treated with 40 units of restriction endonuclease EcoRI to linearize the plasmid. Adenovirus type 5 dl327 DNA (Thimmappaya (1982)) was digested with restriction endonuclease ClaI and the large fragment (approximately 33 kilobase pairs) was purified by sucrose gradient centrifugation. Ten (10)  $\mu$ g of EcoRI treated

Pad/MLP/p53/E1b- and 2.5  $\mu$ g of ClaI treated Ad5 dl327 were mixed and used to transfect approximately  $10^6$  293 cells using the MBS mammalian transfection kit as recommended by the supplier. Eight (8) days following the transfection the 293 cells were split 1 to 3 into fresh media and two days following this adenovirus induced cytopathic effect became evident on the transfected cells. At 13 days post-transfection DNA was prepared from the infected cells using standard procedures (Graham and Prevec (1991)) and analyzed by restriction digestion with restriction endonuclease XhoI. Virus directed expression of p53 was verified following infection of SaoS2 osteosarcoma cells with viral lysate and immunoblotting with an anti-p53 monoclonal antibody designated 1801 (Novocasta Lab. Ltd., U.K.).

## EXPERIMENT NO. II

### MATERIALS AND METHODS

#### Cell Lines

Recombinant adenoviruses were grown and propagated in the human embryonal kidney cell line 293 (ATCC CRL 1573) maintained in DME medium containing 10% defined, supplemented calf serum (Hyclone). Saos-2 cells were maintained in Kaighn's media supplemented with 15% fetal calf serum. HeLa and Hep 3B cells were maintained in DME medium supplemented with 10% fetal calf serum. All other cell lines were grown in Kaighn's media supplemented with 10% fetal calf serum. Saos-2 cells were kindly provided by Dr. Eric Stanbridge. All other cell lines were obtained from ATCC.

#### Construction of Recombinant Adenoviruses

To construct the Ad5/p53 viruses, a 1.4 kb HindIII-SmaI fragment containing the full length cDNA for

p53 (Table I) was isolated from pGEM1-p53-B-T (kindly supplied by Dr. Wen Hwa Lee) and inserted into the multiple cloning site of the expression vector pSP72 (Promega) using standard cloning procedures (Sambrook et al. (1989)). The

5 p53 insert was recovered from this vector following digestion with XhoI-BglII and gel electrophoresis. The p53 coding sequence was then inserted into either pNL3C or pNL3CMV adenovirus gene transfer vectors (kindly provided by Dr. Robert Schneider) which contain the Ad5 5' inverted

10 terminal repeat and viral packaging signals and the E1a enhancer upstream of either the Ad2 major late promoter (MLP) or the human cytomegalovirus immediate early gene promoter (CMV), followed by the tripartite leader CDNA and Ad 5 sequence 3325-5525 bp in a PML2 background. These new

15 constructs replace the E1 region (bp 360-3325) of Ad5 with p53 driven by either the Ad2 MLP (A/M/53) or the human CMV promoter (A/C/53), both followed by the tripartite leader CDNA (see Figure 4). The p53 inserts use the remaining downstream E1b polyadenylation site. Additional MLP and

20 CMV driven p53 recombinants (A/M/N/53, A/C/N/53) were generated which had a further 705 nucleotide deletion of Ad 5 sequence to remove the protein IX (PIX) coding region. As a control, a recombinant adenovirus was generated from the parental PNL3C plasmid without a p53 insert (A/M). A

25 second control consisted of a recombinant adenovirus encoding the beta-galactosidase gene under the control of the CMV promoter (A/C/ $\beta$ -gal). The plasmids were linearized with either Nru I or Eco RI and co-transfected with the large fragment of a Cla I digested Ad 5 d1309 or d1327

30 mutants (Jones and Shenk (1979)) using a Ca/PO<sub>4</sub> transfection kit (Stratagene). Viral plaques were isolated and recombinants identified by both restriction digest analysis and PCR using recombinant specific primers against the tripartite leader CDNA sequence with downstream p53 CDNA

35 sequence. Recombinant virus was further purified by limiting dilution, and virus particles were purified and

titered by standard methods (Graham and van der Erb (1973); Graham and Prevec (1991)).

#### p53 Protein Detection

Saos-2 or Hep 3B cells ( $5 \times 10^5$ ) were infected  
5 with the indicated recombinant adenoviruses for a period of  
24 hours at increasing multiplicities of infection (MOI) of  
plaque forming units of virus/cell. Cells were then washed  
once with PBS and harvested in lysis buffer (50mM Tris-HCl  
Ph 7.5, 250 Mm NaCl, 0.1% NP40, 50mM NaF, 5mM EDTA, 10ug/ml  
10 aprotinin, 10 ug/ml leupeptin, and 1mM PMSF). Cellular  
proteins (approximately 30  $\mu$ g) were separated by 10% SDS-  
PAGE and transferred to nitrocellulose. Membranes were  
incubated with  $\alpha$ -p53 antibody PAb 1801 (Novocastro)  
followed by sheep anti-mouse IgG conjugated with  
15 horseradish peroxidase. p53 protein was visualized by  
chemiluminescence (ECL kit, Amersham) on Kodak XAR-5 film.

#### Measurement of DNA Synthesis Rate

Cells ( $5 \times 10^3$ /well) were plated in 96-well titer  
plates (Costar) and allowed to attach overnight (37°C, 7%  
20 CO<sub>2</sub>). Cells were then infected for 24 hours with purified  
recombinant virus particles at MOIs ranging from 0.3 to 100  
as indicated. Media were changed 24 hours after infection,  
and incubation was continued for a total of 72 hours. <sup>3</sup>H-  
thymidine (Amersham, 1 $\mu$ Ci/well) was added 18 hours prior to  
25 harvest. Cells were harvested on glass fiber filters and  
levels of incorporated radioactivity were measured in a  
beta scintillation counter. <sup>3</sup>H-thymidine incorporation was  
expressed as the mean % (+/- SD) of media control and  
plotted versus the MOI.

30 Tumorigenicity in Nude Mice

Approximately  $2.4 \times 10^8$  Saos-2 cells, plated in T225 flasks, were treated with suspension buffer (1% sucrose in PBS) containing either A/M/N/53 or A/M purified virus at an MOI of 3 or 30. Following an overnight infection, cells were injected subcutaneously into the left and right flanks of BALB/c athymic nude mice (4 mice per group). One flank was injected with the A/M/N/53 treated cells, while the contralateral flank was injected with the control A/M treated cells, each mouse serving as its own control. Animals receiving bilateral injection of buffer treated cells served as additional controls. Tumor dimensions (length, width and height) and body weights were then measured twice per week over an 8 week period. Tumor volumes were estimated for each animal assuming a spherical geometry with radius equal to one-half the average of the measured tumor dimensions.

#### Intra-tumoral RNA Analysis

BALB/c athymic nude mice (approximately 5 weeks of age) were injected subcutaneously with  $1 \times 10^7$  H69 small cell lung carcinoma (SCLC) cells in their right flanks. Tumors were allowed to progress for 32 days until they were approximately 25-50 mm<sup>3</sup>. Mice received peritumoral injections of either A/C/53 or A/C/ $\beta$ -gal recombinant adenovirus ( $2 \times 10^9$  plaque forming units (pfu)) into the subcutaneous space beneath the tumor mass. Tumors were excised from the animals 2 and 7 days post adenovirus treatment and rinsed with PBS. Tumor samples were homogenized, and total RNA was isolated using a TriReagent kit (Molecular Research Center, Inc.). PolyA RNA was isolated using the PolyAtract mRNA Isolation System (Promega), and approximately 10 ng of sample was used for RT-PCR determination of recombinant p53 mRNA expression (Wang et al. (1989)). Primers were designed to amplify sequence between the adenovirus tripartite leader CDNA and

the downstream p53 CDNA, ensuring that only recombinant, and not endogenous p53 would be amplified.

#### p53 Gene Therapy of Established Tumors in Nude Mice

Approximately  $1 \times 10^7$  H69 (SCLC) tumor cells in 200 $\mu$ l volumes were injected subcutaneously into female BALB/c athymic nude mice. Tumors were allowed to develop for 2 weeks, at which point animals were randomized by tumor size (N=5/group). Peritumoral injections of either A/M/N/53 or the control A/M adenovirus ( $2 \times 10^9$  pfu/injection) or buffer alone (1% sucrose in PBS) were administered twice per week for a total of 8 doses/group. Tumor dimensions and body weights were measured twice per week for 7 weeks, and tumor volume was estimated as described previously. Animals were then followed to observe the effect of treatment on mouse survival.

#### RESULTS

##### Construction of Recombinant p53-Adenovirus

p53 adenoviruses were constructed by replacing a portion of the E1a and E1b region of adenovirus Type 5 with p53 CDNA under the control of either the Ad2 MLP (A/M/53) or CMV (A/C/53) promoter (schematized in Figure 4). This E1 substitution severely impairs the ability of the recombinant adenoviruses to replicate, restricting their propagation to 293 cells which supply Ad 5 E1 gene products in trans (Graham et al. (1977)). After identification of p53 recombinant adenovirus by both restriction digest and PCR analysis, the entire p53 CDNA sequence from one of the recombinant adenoviruses (A/M/53) was sequenced to verify that it was free of mutations. Following this, purified preparations of the p53 recombinants were used to infect HeLa cells to assay for the presence of phenotypically wild type adenovirus. HeLa cells, which are non-permissive for

replication of E1-deleted adenovirus, were infected with  $1-4 \times 10^9$  infectious units of recombinant adenovirus, cultured for 3 weeks, and observed for the appearance of cytopathic effect (CPE). Using this assay, recombinant adenovirus replication or wild type contamination was not detected, readily evident by the CPE observed in control cells infected with wild type adenovirus at a level of sensitivity of approximately 1 in  $10^9$ .

#### p53 Protein Expression from Recombinant Adenovirus

To determine if p53 recombinant adenoviruses expressed p53 protein, tumor cell lines which do not express endogenous p53 protein were infected. The human tumor cell lines Saos-2 (osteosarcoma) and Hep 3B (hepatocellular carcinoma) were infected for 24 hours with the p53 recombinant adenoviruses A/M/53 or A/C/53 at MOIs ranging 0.1 to 200 pfu/cell. Western analysis of lysates prepared from infected cells demonstrated a dose-dependent p53 protein expression in both cell types (Figure 5). Both cell lines expressed higher levels of p53 protein following infection with A/C/53 than with A/M/53 (Figure 3). No p53 protein was detected in non-infected cells. Levels of endogenous wild-type p53 are normally quite low, and nearly undetectable by Western analysis of cell extracts (Bartek et al. (1991)). It is clear however that wild-type p53 protein levels are easily detectable after infection with either A/M/53 or A/C/53 at the lower MOIs (Figure 5), suggesting that even low doses of p53 recombinant adenoviruses can produce potentially efficacious levels of p53.

#### p53 Dependent Morphology Changes

The reintroduction of wild-type p53 into the p53-negative osteosarcoma cell line, Saos-2, results in a characteristic enlargement and flattening of these normally

spindle-shaped cells (Chen et al. (1990)). Subconfluent Saos-2 cells ( $1 \times 10^5$  cells/10cm plate) were infected at an MOI of 50 with either the A/C/53 or control A/M virus, and incubated at 37°C for 72 hours until uninfected control plates were confluent. At this point, the expected morphological change was evident in the A/C/53 treated plate (Figure 6, panel C) but not in uninfected (Figure 6, panel A) or control virus-infected plates (Figure 6, panel B). This effect was not a function of cell density because a control plate initially seeded at lower density retained normal morphology at 72 hours when its confluence approximated that of the A/C/53 treated plate. Previous results had demonstrated a high level of p53 protein expression at an MOI of 50 in Saos-2 cells (Figure 5A), and these results provided evidence that the p53 protein expressed by these recombinant adenoviruses was biologically active.

#### p53 Inhibition of Cellular DNA Synthesis

To further test the activity of the p53 recombinant adenoviruses, their ability to inhibit proliferation of human tumor cells was assayed as measured by the uptake of  $^3\text{H}$ -thymidine. It has previously been shown that introduction of wild-type p53 into cells which do not express endogenous wild-type p53 can arrest the cells at the  $G_1/S$  transition, leading to inhibition of uptake of labeled thymidine into newly synthesized DNA (Baker et al. (1990); Mercer et al. (1990); Diller et al. (1990)). A variety of p53-deficient tumor cell lines were infected with either A/M/N/53, A/C/N/53 or a non-p53 expressing control recombinant adenovirus (A/M). A strong, dose-dependent inhibition of DNA synthesis by both the A/M/N/53 and A/C/N/53 recombinants in 7 out of the 9 different tumor cell lines tested (Figure 7) was observed. Both constructs were able to inhibit DNA synthesis in these human tumor cells, regardless of whether they expressed mutant p53 or



failed to express p53 protein. It also was found that in this assay, the A/C/N/53 construct was consistently more potent than the A/M/N/53. In saos-2 (osteosarcoma) and MDA-MB468 (breast cancer) cells, nearly 100% inhibition of DNA synthesis was achieved with the A/C/N/53 construct at an MOI as low as 10. At doses where inhibition by the control adenovirus in only 10-30%, a 50-100% reduction in DNA synthesis using either p53 recombinant adenovirus was observed. In contrast, no significant p53-specific effect was observed with either construct as compared to control virus in HEP G2 cells (hepatocarcinoma cell line expressing endogenous wild-type p53, Bressac et al. (1990)), nor in the K562 (p53 null) leukemic cell line.

2025 RELEASE UNDER E.O. 14176

### Tumorigenicity in Nude Mice

In a more stringent test of function for the p53 recombinant adenoviruses, tumor cells were infected *ex vivo* and then injected the cells into nude mice to assess the ability of the recombinants to suppress tumor growth *in vivo*. Saos-2 cells infected with A/M/N/53 or control A/M virus at a MOI of 3 or 30, were injected into opposite flanks of nude mice. Tumor sizes were then measured twice a week over an 8 week period. At the MOI of 30, no tumor growth was observed in the p53-treated flanks in any of the animals, while the control treated tumors continued to grow (Figure 8). The progressive enlargement of the control virus treated tumors were similar to that observed in the buffer treated control animals. A clear difference in tumor growth between the control adenovirus and the p53 recombinant at the MOI of 3, although tumors from 2 out of the 4 p53-treated mice did start to show some growth after approximately 6 weeks. Thus, the A/M/N/53 recombinant adenovirus is able to mediate p53-specific tumor suppression in an *in vivo* environment.

### *In Vivo* Expression of Ad/p53

Although *ex vivo* treatment of cancer cells and subsequent injection into animals provided a critical test of tumor suppression, a more clinically relevant experiment is to determine if injected p53 recombinant adenovirus could infect and express p53 in established tumors *in vivo*. To address this, H69 (SCLC, p53<sup>null</sup>) cells were injected subcutaneously into nude mice, and tumors were allowed to develop for 32 days. At this time, a single injection of 2 x 10<sup>9</sup> pfu of either A/C/53 or A/C/ $\beta$ -gal adenovirus was injected into the peritumoral space surrounding the tumor. Tumors were then excised at either Day 2 or Day 7 following the adenovirus injection, and polyA RNA was isolated from each tumor. RT-PCR, using recombinant-p53 specific

primers, was then used to detect p53 mRNA in the p53 treated tumors (Figure 9, lanes 1,2,4,5). No p53 signal was evident from the tumors excised from the  $\beta$ -gal treated animals (Figure 9, lanes 3 and 6). Amplification with actin primers served as a control for the RT-PCR reaction (Figure 9, lanes 7-9), while a plasmid containing the recombinant-p53 sequence served as a positive control for the recombinant-p53 specific band (Figure 9, lane 10). This experiment demonstrates that a p53 recombinant adenovirus can specifically direct expression of p53 mRNA within established tumors following a single injection into the peritumoral space. It also shows *in vivo* viral persistence for at least one week following infection with a p53 recombinant adenovirus.

#### 15 *In Vivo* Efficacy

To address the feasibility of gene therapy of established tumors, a tumor-bearing nude mouse model was used. H69 cells were injected into the subcutaneous space on the right flank of mice, and tumors were allowed to grow for 2 weeks. Mice then received peritumoral injections of buffer or recombinant virus twice weekly for a total of 8 doses. In the mice treated with buffer or control A/M virus, tumors continued to grow rapidly throughout the treatment, whereas those treated with the A/M/N/53 virus grew at a greatly reduced rate (Figure 10A). After cessation of injections, the control treated tumors continued to grow while the p53 treated tumors showed little or no growth for at least one week in the absence of any additional supply of exogenous p53 (Figure 10A). Although control animals treated with buffer alone had accelerated tumor growth as compared to either virus treated group, no significant difference in body weight was found between the three groups during the treatment period. Tumor ulceration in some animals limited the relevance of tumor size measurements after day 42. However, continued

monitoring of the animals to determine survival time demonstrated a survival advantage for the p53-treated animals (Figure 10B). The last of the control adenovirus treated animals died on day 83, while buffer alone treated controls had all expired by day 56. In contrast, all 5 animals treated with the A/M/N/53 continue to survive (day 130 after cell inoculation) (Figure 10B). Together, this data establish a p53-specific effect on both tumor growth and survival time in animals with established p53-deficient tumors.

#### Adenovirus Vectors Expressing p53

Recombinant human adenovirus vectors which are capable of expressing high levels of wild-type p53 protein in a dose dependent manner were constructed. Each vector contains deletions in the E1a and E1b regions which render the virus replication deficient (Challberg and Kelly (1979); Horowitz, (1991)). Of further significance is that these deletions include those sequences encoding the E1b 19 and 55 kd protein. The 19 kd protein is reported to be involved in inhibiting apoptosis (White et al. (1992); Rao et al. (1992)), whereas the 55 kd protein is able to bind wild-type p53 protein (Sarnow et al. (1982); Heuvel et al. (1990)). By deleting these adenoviral sequences, potential inhibitors of p53 function were removed through direct binding to p53 or potential inhibition of p53 mediated apoptosis. Additional constructs were made which have had the remaining 3' E1b sequence, including all protein IX coding sequence, deleted as well. Although this has been reported to reduce the packaging size capacity of adenovirus to approximately 3 kb less than wild-type virus (Ghosh-Choudhury et al. (1987)), these constructs are also deleted in the E3 region so that the A/M/N/53 and A/C/N/53 constructs are well within this size range. By deleting the pIX region, adenoviral sequences homologous to those contained in 293 cells are reduced to approximately 300

base pairs, decreasing the chances of regenerating replication-competent, wild-type adenovirus through recombination. Constructs lacking pIX coding sequence appear to have equal efficacy to those with pIX.

5 p53/Adenovirus Efficacy In Vitro

10 In concordance with a strong dose dependency for expression of p53 protein in infected cells, a dose-dependent, p53-specific inhibition of tumor cell growth was demonstrated. Cell division, was inhibited and demonstrated by the inhibition of DNA synthesis, in a wide variety of tumor cell types known to lack wild-type p53 protein expression. Bacchetti and Graham (1993) recently reported p53 specific inhibition of DNA synthesis in the ovarian carcinoma cell line SKOV-3 by a p53 recombinant adenovirus in similar experiments. In addition to ovarian carcinoma, additional human tumor cell lines were demonstrated, representative of clinically important human cancers and including lines over-expressing mutant p53 protein, can also be growth inhibited by the p53 recombinants of this invention. At MOIs where the A/C/N/53 recombinant is 90-100% effective in inhibiting DNA synthesis in these tumor types, control adenovirus mediated suppression is less than 20%.

25 Although Feinstein et al. (1992) reported that re-introduction of wild-type p53 could induce differentiation and increase the proportion of cells in G<sub>1</sub> versus S+G<sub>2</sub> for leukemic K562 cells, no p53 specific effect was found in this line. Horvath and Weber (1988) have reported that human peripheral blood lymphocytes are highly nonpermissive to adenovirus infection. In separate experiments, the recombinant significantly infected the non-responding K562 cells with recombinant A/C/β-gal adenovirus, while other cell lines, including the control Hep G2 line and those showing a strong p53 effect, were

readily infectable. Thus, at least part of the variability of efficacy would appear to be due to variability of infection, although other factors may be involved as well.

The results observed with the A/M/N/53 virus in Figure 8 demonstrates that complete suppression is possible in an *in vivo* environment. The resumption of tumor growth in 2 out of 4, p53 treated animals at the lower MOI most likely resulted from a small percentage of cells not initially infected with the p53 recombinant at this dose. The complete suppression seen with A/M/N/53 at the higher dose, however, shows that the ability of tumor growth to recover can be overcome.

#### p53/Adenovirus *In Vivo* Efficacy

Work presented here and by other groups (Chen et al. (1990); Takahashi et al. (1992)) have shown that human tumor cells lacking expression of wild-type p53 can be treated *ex vivo* with p53 and result in suppression of tumor growth when the treated cells are transferred into an animal model. Applicants present the first evidence of tumor suppressor gene therapy of an *in vivo* established tumor, resulting in both suppression of tumor growth and increased survival time. In Applicants' system, delivery to tumor cells did not rely on direct injection into the tumor mass. Rather, p53 recombinant adenovirus was injected into the peritumoral space, and p53 mRNA expression was detected within the tumor. p53 expressed by the recombinants was functional and strongly suppressed tumor growth as compared to that of control, non-p53 expressing adenovirus treated tumors. However, both p53 and control virus treated tumor groups showed tumor suppression as compared to buffer treated controls. It has been demonstrated that local expression of tumor necrosis factor (TNF), interferon- $\gamma$ , interleukin (IL)-2, IL-4 or IL-7 can lead to T-cell independent transient tumor

suppression in nude mice (Hoch et al. (1992)). Exposure of monocytes to adenovirus virions are also weak inducers of IFN- $\alpha/\beta$  (reviewed in Gooding and Wold (1990)). Therefore, it is not surprising that some tumor suppression in nude mice was observed even with the control adenovirus. This virus mediated tumor suppression was not observed in the ex vivo control virus treated Saos-2 tumor cells described earlier. The p53-specific in vivo tumor suppression was dramatically demonstrated by continued monitoring of the animals in Figure 10. The survival time of the p53-treated mice was significantly increased, with 5 out of 5 animals still alive more than 130 days after cell inoculation compared to 0 out of 5 adenovirus control treated animals. The surviving animals still exhibit growing tumors which may reflect cells not initially infected with the p53 recombinant adenovirus. Higher or more frequent dosing schedules may address this. In addition, promoter shutoff (Palmer et al. (1991)) or additional mutations may have rendered these cells resistant to the p53 recombinant adenovirus treatment. For example, mutations in the recently described WAF1 gene, a gene induced by wild-type p53 which subsequently inhibits progression of the cell cycle into S phase, (El-Deiry et al. (1993); Hunter (1993)) could result in a p53-resistant tumor.

25

**EXPERIMENT NO. III**

This Example shows the use of suicide genes and tissue specific expression of such genes in the gene therapy methods described herein. Hepatocellular carcinoma was chosen as the target because it is one of the most common human malignancies affecting man, causing an estimated 1,250,000 deaths per year world-wide. The incidence of this cancer is very high in Southeast Asia and Africa where it is associated with Hepatitis B and C infection and exposure to aflatoxin. Surgery is currently the only treatment which offers the potential for curing

HCC, although less than 20% of patients are considered candidates for resection (Ravoet C. et al., 1993). However, tumors other than hepatocellular carcinoma are equally applicable to the methods of reducing their proliferation described herein.

#### CELL LINES

All cell lines but for the HLF cell line were obtained from the American Type Tissue Culture Collection (ATCC) 12301 Parklawn Drive, Rockville Maryland. ATCC accession numbers are noted in parenthesis. The human embryonal kidney cell line 293 (CRL 1573) was used to generate and propagate the recombinant adenoviruses described herein. They were maintained in DME medium containing 10% defined, supplemented calf serum (Hyclone). The hepatocellular carcinoma cell lines Hep 3B (HB 8064), Hep G2 (HB 8065), and HLF were maintained in DME/F12 medium supplemented with 10% fetal bovine serum, as were the breast carcinoma cell lines MDA-MB468 (HTB 132) and BT-549 (HTB 122). Chang liver cells (CCL 13) were grown in MEM medium supplemented with 10% fetal bovine serum. The HLF cell line was obtained from Drs. T. Morsaki and H. Kitsuki at the Kyushu University School of Medicine in Japan.

#### RECOMBINANT VIRUS CONSTRUCTION

Two adenoviral expression vectors designated herein as ACNTK and AANTK and devoid of protein IX function (depicted in Figure 11) are capable of directing expression of the TK suicide gene within tumor cells. A third adenovirus expression vector designated AANCAT was constructed to further demonstrate the feasibility of specifically targeting gene expression to specific cell types using adenoviral vectors. These adenoviral constructs were assembled as depicted in Figures 11 and 12 and are derivatives of those previously described for the expression of tumor suppressor genes.



For expression of the foreign gene, expression cassettes have been inserted that utilize either the human cytomegalovirus immediate early promoter/enhancer (CMV) (Boshart, M. et al., 1985) or the human alpha-fetoprotein (AFP) enhancer/promoter (Watanabe, K. et al., 1987; Nakabayashi, H. et al., 1989) to direct transcription of the TK gene or the chloramphenicol acetyltransferase gene (CAT). The CMV enhancer promoter is capable of directing robust gene expression in a wide variety of cell types while the AFP enhancer/promoter construct restricts expression to hepatocellular carcinoma cells (HCC) which express AFP in about 70-80% of the HCC patient population. In the construct utilizing the CMV promoter/enhancer, the adenovirus type 2 tripartite leader sequence also was inserted to enhance translation of the TK transcript (Berkner, K.L. and Sharp, 1985). In addition to the E1 deletion, both adenovirus vectors are additionally deleted for 1.9 kilobases (kb) of DNA in the viral E3 region. The DNA deleted in the E3 region is non-essential for virus propagation and its deletion increases the insert capacity of the recombinant virus for foreign DNA by an equivalent amount (1.9kb) (Graham and Prevec, 1991).

To demonstrate the specificity of the AFP promoter/enhancer, the virus AANCAT also was constructed where the marker gene chloramphenicol acetyltransferase (CAT) is under the control of the AFP enhancer/promoter. In the ACNTK viral construct, the Ad2 tripartite leader sequence was placed between the CMV promoter/enhancer and the TK gene. The tripartite leader has been reported to enhance translation of linked genes. The E1 substitution impairs the ability of the recombinant viruses to replicate, restricting their propagation to 293 cells which supply the Ad5 E1 gene products in trans (Graham et al., 1977).

Adenoviral Vector ACNTK: The plasmid pMLBKTK in E. coli HB101 (from ATCC #39369) was used as the source of the herpes simplex virus (HSV-1) thymidine kinase (TK) gene. TK was excised from this plasmid as a 1.7 kb gene  
5 fragment by digestion with the restriction enzymes Bgl II and Pvu II and subcloned into the compatible Bam HI, EcoR V restriction sites of plasmid pSP72 (Promega) using standard cloning techniques (Sambrook et al., 1989). The TK insert was then isolated as a 1.7 kb fragment from this  
10 vector by digestion with Xba I and Bgl II and cloned into Xba I, BamHI digested plasmid pACN (Wills et al. 1994). Twenty (20)  $\mu$ g of this plasmid designated pACNTK were linearized with Eco RI and cotransfected into 293 cells (ATCC CRL 1573) with 5  $\mu$ g of Cla I digested ACBGL (Wills et  
15 al., 1994 supra) using a CaPO<sub>4</sub> transfection kit (Stratagene, San Diego, California). Viral plaques were isolated and recombinants, designated ACNTK, were identified by restriction digest analysis of isolated DNA with Xho I and BsiWI. Positive recombinants were further purified by  
20 limiting dilution and expanded and titered by standard methods (Graham and Prevec, 1991).

Adenoviral Vector AANTK: The  $\alpha$ -fetoprotein promoter (AFP-P) and enhancer (AFP-E) were cloned from a human genomic DNA (Clontech) using PCR amplification with  
25 primers containing restriction sites at their ends. The primers used to isolate the 210 bp AFP-E contained a Nhe I restriction site on the 5' primer and an Xba I, Xho I, Kpn I linker on the 3' primer. The 5' primer sequence was 5'-CGC GCT AGC TCT GCC CCA AAG AGC T-3. The 5' primer  
30 sequence was 5'-CGC GGT ACC CTC GAG TCT AGA TAT TGC CAG TGG TGG AAG-3'. The primers used to isolate the 1763 bp AFE fragment contained a Not I restriction site on the 5' primer and a Xba I site on the 3' primer. The 5' primer sequence was 5'-CGT GCG GCC GCT GGA GGA CTT TGA GGA TGT CTG  
35 TC-3'. The 3' primer sequence was 5'-CGC TCT AGA GAG ACC AGT TAG GAA GTT TTC GCA-3'. For PCR amplification, the DNA

was denatured at 97° for 7 minutes, followed by 5 cycles of amplification at 97°, 1 minute, 53°, 1 minute, 72°, 2 minutes, and a final 72°, 10 minute extension. The amplified AFE was digested with Not I and Xba I and inserted into the Not I, Xba I sites of a plasmid vector (pA/ITR/B) containing adenovirus type 5 sequences 1-350 and 3330 - 5790 separated by a polylinker containing Not I, Xho I, Xba I, Hind III, Kpn I, Bam HI, Nco I, Sma I, and Bgl II sites. The amplified AFP-E was digested with Nhe I and Kpn I and inserted into the AFP-E containing construct described above which had been digested with Xba I and Kpn I. This new construct was then further digested with Xba I and NgoMI to remove adenoviral sequences 3330 - 5780, which were subsequently replaced with an Xba I, NgoMI restriction fragment of plasmid pACN containing nucleotides 4021 - 10457 of adenovirus type 2 to construct the plasmid pAAN containing both the  $\alpha$ -fetoprotein enhancer and promoter. This construct was then digested with Eco RI and Xba I to isolate a 2.3 kb fragment containing the Ad5 inverted terminal repeat, the AFP-E and the AFP-P which was subsequently ligated with the 8.55 kb fragment of Eco RI, Xba I digested pACNTK described above to generate pAANTK where the TK gene is driven by the  $\alpha$ -fetoprotein enhancer and promoter in an adenovirus background. This plasmid was then linearized with Eco RI and cotransfected with the large fragment of Cla I digested ALBGL as above and recombinants, designated AANTK, were isolated and purified as described above.

Adenoviral Vector AANCAT: The chloramphenicol acetyltransferase (CAT) gene was isolated from the pCAT-Basic Vector (Promega Corporation) by an Xba I, Bam HI digest. This 1.64 kb fragment was ligated into Xba I, Bam HI digested pAAN (described above) to create pAANCAT. This plasmid was then linearized with Eco RI and cotransfected with the large fragment of Cla I digested rA/C/ $\beta$ -gal to create AANCAT.

REPORTER GENE EXPRESSION:  $\beta$ -GALACTOSIDASE EXPRESSION:

Cells were plated at  $1 \times 10^5$  cells/well in a 24-well tissue culture plate (Costar) and allowed to adhere overnight (37°C, 7% CO<sub>2</sub>). Overnight infections of ACBGL were performed at a multiplicity of infection (MOI) of 30. After 24 hours, cells were fixed with 3.7% Formaldehyde; PBS, and stained with 1mg/ml Xgal reagent (USB). The data was scored (+, ++, +++) by estimating the percentage of positively stained cells at each MOI. [+=1-33%, ++=33-67% and +++=>67%]

## REPORTER GENE EXPRESSION: CAT EXPRESSION:

Two (2)  $\times 10^6$  cells (Hep G2, Hep 3B, HLF, Chang, and MDA-MB468) were seeded onto 10 cm plates in triplicate and incubated overnight (37°C, 7% CO<sub>2</sub>). Each plate was then infected with either AANCAT at an MOI = 30 or 100 or uninfected and allowed to incubate for 3 days. The cells were then trypsinized and washed with PBS and resuspended in 100  $\mu$ l of 0.25 M Tris pH 7.8. The samples were frozen and thawed 3 times, and the supernatant was transferred to new tubes and incubated at 60°C for 10 minutes. The samples were then spun at 4°C for 5 minutes, and the supernatants assayed for protein concentration using a Bradford assay (Bio-Rad Protein Assay Kit). Samples were adjusted to equal protein concentrations to a final volume of 75  $\mu$ l using 0.25 M Tris, 25  $\mu$ l of 4mM acetyl CoA and 1  $\mu$ l of <sup>14</sup>C-Chloramphenicol and incubated overnight at 37°C. 500  $\mu$ l of ethyl acetate is added to each sample and mixed by vortexing, followed by centrifugation for 5 minutes at room temperature. The upper phase is then transferred to a new tube and the ethyl acetate is evaporated by centrifugation under vacuum. The reaction products are then redissolved in 25  $\mu$ l of ethyl acetate and spotted onto a thin layer chromatography (TLC) plate and the plate is then placed in a pre-equilibrated TLC chamber (95%

chloroform, 5% methanol). The solvent is then allowed to migrate to the top of the plate, the plate is then dried and exposed to X-ray film.

#### CELLULAR PROLIFERATION: $^3\text{H}$ -THYMIDINE INCORPORATION

5 Cells were plated at  $5 \times 10^3$  cells/well in a 96-well micro-titer plate (Costar) and allowed to incubate overnight (37C, 7%  $\text{CO}_2$ ). Serially diluted ACN, ACNTK or AATK virus in DMEM; 15% FBS; 1% glutamine was used to  
10 infect cells at an infection multiplicity of 30 for an overnight duration at which point cells were dosed in triplicate with ganciclovir (Cytovene) at log intervals between 0.001 and 100  $\mu\text{M}$  (micro molar). 1  $\mu\text{Ci}$   $^3\text{H}$ -thymidine (Amersham) was added to each well 12-18 hours before  
15 harvesting. At 72 hours-post infection cells were harvested onto glass-fiber filters and incorporated  $^3\text{H}$ -thymidine was counted using liquid scintillation (TopCount, Packard). Results are plotted as percent of untreated control proliferation and tabulated as the effective dose ( $\text{ED}_{50} \pm \text{SD}$ ) for a 50 percent reduction in proliferation over  
20 media controls.  $\text{ED}_{50}$  values were estimated by fitting a logistic equation to the dose response data.

#### CYTOTOXICITY: LDH RELEASE

Cells (HLF, human HCC) were plated, infected with ACN or ACNTK and treated with ganciclovir as described  
25 for the proliferation assay. At 72 hours post-ganciclovir administration, cells were spun, the supernatant was removed. The levels of lactate dehydrogenase measured colorimetrically (Promega, Cytotox 96<sup>TM</sup>). Mean ( $\pm$  S.D.) LDH release is plotted versus M.O.I.

#### 30 IN VIVO THERAPY

Human hepatocellular carcinoma cells (Hep 3B) were injected subcutaneously into ten female (10) athymic nu/nu mice (Simonsen Laboratories, Gilroy, CA). Each animal received approximately  $1 \times 10^7$  cells in the left flank. Tumors were allowed to grow for 27 days before randomizing mice by tumor size. Mice were treated with intratumoral and peritumoral injections of ACNTK or the control virus ACN ( $1 \times 10^9$  iu in  $100 \mu\text{l}$ ) every other day for a total of three doses. Starting 24 hours after the initial dose of adenovirus, the mice were dosed intraperitoneally with ganciclovir (Cytovene 100 mg/kg) daily for a total of 10 days. Mice were monitored for tumor size and body weight twice weekly. Measurements on tumors were made in three dimensions using vernier calipers and volumes were calculated using the formula  $\frac{4}{3} \pi r^3$ , where r is one-half the average tumor dimension.

#### RESULTS

The recombinant adenoviruses were used to infect three HCC cell lines (HLF, Hep3B and Hep-G2). One human liver cell line (Chang) and two breast cancer cell lines were used as controls (MDAMB468 and BT549). To demonstrate the specificity of the AFP promoter/enhancer, the virus AANCAT was constructed. This virus was used to infect cells that either do (Hep 3B, HepG2) or do not (HLE, Chang, MDAMB468) express the HCC tumor marker alpha-fetoprotein (AFP). As shown in Figure 13, AANCAT directs expression of the CAT marker gene only in those HCC cells which are capable of expressing AFP (Figure 13).

The efficacy of ACNTK and AANTK for the treatment of HCC was assessed using a  $^3\text{H}$ -thymidine incorporation assay to measure the effect of the combination of HSV-TK expression and ganciclovir treatment upon cellular proliferation. The cell lines were infected with either ACNTK or AANTK or the control virus ACN (Wills et al., 1994

supra), which does not direct expression of HSV-TK, and then treated with increasing concentrations of ganciclovir. The effect of this treatment was assessed as a function of increasing concentrations of ganciclovir, and the concentration of ganciclovir required to inhibit <sup>3</sup>H-thymidine incorporated by 50% was determined (ED<sub>50</sub>). Additionally, a relative measure of adenovirus - mediated gene transfer and expression of each cell line was determined using a control virus which directs expression of the marker gene beta-galactosidase. The data presented in Figure 14 and Table 1 below show that the ACNTK virus/ganciclovir combination treatment was capable of inhibiting cellular proliferation in all cell lines examined as compared with the control adenovirus ACN in combination with ganciclovir. In contrast, the AANTK viral vector was only effective in those HCC cell lines which have been demonstrated to express  $\alpha$ -fetoprotein. In addition, the AANTK/GCV combination was more effective when the cells were plated at high densities.

TABLE 1

Cell Line	aFP	$\beta$ -gal Expression	ACN	ED50 ACNTK	AANTK
MDAMB468	-	+++	>100	2	>100
BT549	-	+++	>100	<0.3	>100
HLF	-	+++	>100	0.8	>100
CHANG	-	+++	>100	22	>100
HEP-3B	-	+	80	8	8
HEP-G2 LOW	+	++	90	2	35
HEP-G2 HIGH	+	++	89	0.5	4

Nude mice bearing Hep3B tumors (N=5/group) were treated intratumorally and peritumorally with equivalent

5 doses of ACNTK or ACN control. Twenty-four hours after the first administration of recombinant adenovirus, daily treatment of ganciclovir was initiated in all mice. Tumor dimensions from each animal were measured twice weekly via  
10 calipers, and average tumor sizes are plotted in Figure 16. Average tumor size at day 58 was smaller in the ACNTK-treated animals but the difference did not reach statistical significance ( $p < 0.09$ , unpaired t-test). These data support a specific effect of ACNTK on tumor growth in vivo. No significant differences in average body weight were detected between the groups.

15 Although the invention has been described with reference to the above embodiments, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the claims that follow.

0055570-10289  
268207-02585580



## REFERENCES

- AIELLO, L. et al. (1979) Virology 94:460-469.
- AMERICAN CANCER SOCIETY. (1993) Cancer Facts and Figures.
- AULITZKY et al. (1991) Eur. J. Cancer 27(4):462-467.
- 5 AUSTIN, E.A. and HUBER, B.E. (1993) Mol. Pharmaceutical  
43:380-387.
- BACCHETTI, S. AND GRAHAM, F. (1993) International Journal  
of Oncology 3:781-788.
- BAKER S.J., MARKOWITZ, S., FEARON E.R., WILLSON, J.K.V.,  
10 AND VOGELSTEIN, B. (1990) Science 249:912-915.
- BARTEK, J., BARTKOVA, J., VOJTESEK, B., STASKOVA, Z.,  
LUKAS, J., REJTHAR, A., KOVARIK, J., MIDGLEY, C.A., GANNON,  
J.V., AND LANE, D.P. (1991) Oncogene 6:1699-1703.
- BERKNER, K.L. and SHARP (1985) Nucleic Acids Res 13:841-  
15 857.
- BOSHART, M. et al. (1985) Cell 41:521-530.
- BRESSAC, B., GALVIN, K.M., LIANG, T.J., ISSELBACHER, K.J.,  
WANDS, J.R., AND OZTURK, M. (1990) Proc. Natl. Acad. Sci.  
USA 87:1973-1977.
- 20 CARUSO M. et al. (1993) Proc. Natl. Acad. Sci. USA 90:7024-  
7028.
- CHALLBERG, M.D., KELLY, T.J. (1979) Biochemistry 76:655-  
659.
- CHEN P.L., CHEN Y., BOOKSTEIN R., AND LEE W.H. (1990)  
25 Science 250:1576-1580.

2025 RELEASE UNDER E.O. 14176

CHEN, Y., CHEN, P.L., ARNAIZ, N., GOODRICH, D., AND LEE, W.H. (1991) *Oncogene* 6:1799-1805.

CHENG, J.L., YEE, J.K., YEARGIN, J., FRIEDMANN, T., AND HAAS, M. (1992) *Cancer Research* 52:222-226.

- 5 COLBY, W.W. AND SHENK, T. J. (1981) *Virology* 39:977-980.

CULVER ET AL. (1991) *P.N.A.S. (U.S.A.)* 88:3155-3159.

CULVER, K.W. et al. (1992) *Science* 256:1550-1552.

DEMETRI et al. (1989) *J. Clin. Oncol.* 7(10):1545-1553.

DILLER, L., et al. (1990) *Mol. Cell. Biology* 10:5772-5781.

- 10 EL-DEIRY, W.S., et al. (1993) *Cell* 75:817-825.

EZZIDINE, Z.D. et al. (1991) *The New Biologist* 3:608-614.

FEINSTEIN, E., GALE, R.P., REED, J., AND CANAANI, E. (1992) *Oncogene* 7:1853-1857.

- 15 GHOSH-CHOUDHURY, G., HAJ-AHMAD, Y., AND GRAHAM, F.L. (1987) *EMBO Journal* 6:1733-1739.

GOODING, L.R., AND WOLD, W.S.M. (1990) *Crit. Rev. Immunol.* 10:53-71.

GRAHAM F.L., AND VAN DER ERB A.J. (1973) *Virology* 52:456-467.

- 20 GRAHAM, F.L. AND PREVEC, L. (1992) Vaccines: New Approaches to Immunological Problems. R.W. Ellis (ed), Butterworth-Heinemann, Boston. pp. 363-390.

202507-02585580

GRAHAM, F.L., SMILEY, J., RUSSELL, W.C. AND NAIRN, R.  
(1977) J. Gen. Virol. 36:59-74.

GRAHAM F.L. AND PREVEC L. (1991) Manipulation of  
adenovirus vectors. In: Methods in Molecular Biology. Vol  
5 7: Gene Transfer and Expression Protocols. Murray E.J.  
(ed.) The Humana Press Inc., Clifton N.J., Vol 7:109-128.

HEUVEL, S.J.L., LAAR, T., KAST, W.M., MELIEF, C.J.M.,  
ZANTEMA, A., AND VAN DER EB, A.J. (1990) EMBO Journal  
9:2621-2629.

10 HOCK, H., DORSCH, M., KUZENDORF, U., QIN, Z., DIAMANTSTEIN,  
T., AND BLANKENSTEIN, T. (1992) Proc. Natl. Acad. Sci. USA  
90:2774-2778.

HOLLSTEIN, M., SIDRANSKY, D., VOGELSTEIN, B., AND HARRIS,  
C. (1991) Science 253:49-53.

15 HOROWITZ, M.S. (1991) Adenoviridae and their replication.  
In Fields Virology. B.N. Fields, ed. (Raven Press, New  
York) pp. 1679-1721.

HORVATH, J., AND WEBER, J.M. (1988) J. Virol. 62:341-345.

HUANG et al. (1991) Nature 350:160-162.

20 HUBER, B.E. et al. (1991) Proc. Natl. Acad. Sci. USA  
88:8039-8043.

HUNTER, T. (1993) Cell 75:839-841.

JONES, N. AND SHENK, T. (1979) Cell 17:683-689.

KAMB et al. (1994) Science 264:436-440.

2025 RELEASE UNDER E.O. 14176

KEURBITZ, S.J., PLUNKETT, B.S., WALSH, W.V., AND KASTAN, M.B. (1992) Proc. Natl. Acad. Sci. USA **89**: 7491-7495.

KREIGLER, M. Gene Transfer and Expression: A Laboratory Manual, W.H. Freeman and Company, New York (1990).

- 5 LANDMANN et al. (1992) J. Interferon Res. **12**(2):103-111.

LANE, D.P. (1992) Nature **358**:15-16.

LANTZ et al. (1990) Cytokine **2**(6):402-406.

- LARRICK, J.W. and BURCK, K.L. Gene Therapy: Application of Molecular Biology, Elsevier Science Publishing Co., Inc.  
10 New York, New York (1991).

LEE et al. (1987) Science **235**:1394-1399.

LEMAISTRE et al. (1991) Lancet **337**:1124-1125.

LEMARCHAND, P., et al. (1992) Proc. Natl. Acad. Sci. USA **89**:6482-6486.

- 15 LEVINE, A.J. (1993) The Tumor Suppressor Genes. Annu. Rev. Biochem. 1993. **62**:623-651.

LOWE S.W., SCHMITT, E.M., SMITH, S.W., OSBORNE, B.A., AND JACKS, J. (1993) Nature **362**:847-852.

- LOWE, S.W., RULEY, H.E., JACKS, T., AND HOUSMAN, D.E.  
20 (1993) Cell **74**:957-967.

MARTIN (1975) In: Remington's Pharm. Sci., 15th Ed. (Mack Publ. Co., Easton).

MERCER, W.E., et al. (1990) Proc. Natl. Acad. Sci. USA **87**:6166-6170.

2025 RELEASE UNDER E.O. 14176

NAKABAYASHI, H. et al. (1989) The Journal of Biological Chemistry **264**:266-271.

PALMER, T.D., ROSMAN, G.J., OSBORNE, W.R., AND MILLER, A.D. (1991) Proc. Natl. Acad. Sci USA **88**:1330-1334.

- 5 RAO, L., DEBBAS, M., SABBATINI, P., HOCKENBERY, D., KORSMEYER, S., AND WHITE, E. (1992) Proc. Natl. Acad. Sci. USA **89**:7742-7746.

RAVOET C. et al. (1993) Journal of Surgical Oncology Supplement **3**:104-111.

- 10 RICH, D.P., et al. (1993) Human Gene Therapy **4**:460-476.

ROSENFELD, M.A., et al. (1992) Cell **68**:143-155.

SAMBROOK J., FRITSCH E.F., AND MANIATIS T. (1989). Molecular Cloning: A Laboratory Manual. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor).

- 15 SARNOW, P., HO, Y.S., WILLIAMS, J., AND LEVINE, A.J. (1982) Cell **28**:387-394.

SHAW, P., BOVEY, R., TARDY, S., SAHLI, R., SORDAT, B., AND COSTA, J. (1992) Proc. Natl. Acad. Sci. USA **89**:4495-4499.

SIEGFRIED, W. (1993) Exp. Clin. Endocrinol. **101**:7-11.

- 20 SORSCHER, E.J. et al. (1994) Gene Therapy **1**:233-238.

SPECTOR, D.J. (1983) Virology **130**:533-538.

STEWART, P.L. et al. (1993) EMBO Journal **12**:2589-2599.

2025-04-04 10:55:50

STRAUS. S.E. (1984) Adenovirus infections in humans. In: The Adenoviruses, Ginsberg HS, ed. New York: Plenum Press, 451-496.

SUPERSAXO et al. (1988) Pharm. Res. 5(8):472-476.

5 TAKAHASHI, T., et al. (1989) Science 246: 491-494.

TAKAHASHI, T., et al. (1992) Cancer Research 52:2340-2343.

THIMMAPPAYA, B. et al. (1982) Cell 31:543-551.

WANG, A.M., DOYLE, M.V., AND MARK, D.F. (1989) Proc. Natl. Acad. Sci USA 86:9717-9721.

10 WATANABLE, K. et al. (1987) The Journal of Biological Chemistry 262:4812-4818.

WHITE, E., et al. (1992) Mol. Cell. Biol. 12:2570-2580.

WILLS, K.N. et al. (1994) Hum. Gen. Ther. 5:1079-1088.

YONISH-ROUACH, E., et al. (1991) Nature 352:345-347.

2025 RELEASE UNDER E.O. 14176

What is claimed is:

1. A recombinant adenovirus expression vector comprising a partial or total deletion of a protein IX DNA and a gene encoding a foreign protein.

2. The recombinant adenovirus expression vector of claim 1, wherein the deletion of the protein IX gene sequence extends from about 3500 bp from the 5' viral termini to about 4000 bp from the 5' viral termini.

3. The recombinant adenovirus expression vector of claim 2 further comprising deletion of a non-essential DNA sequence in adenovirus early region 3 and/or early region 4.

4. The recombinant adenovirus expression vector of claim 2 further comprising deletion of a DNA sequences designated adenovirus E1a and E1b.

5. The recombinant adenovirus expression vector of claim 2 further comprising deletion of early region 3 and/or 4 and DNA sequences designated adenovirus E1a and E1b.

6. The recombinant adenovirus expression vector of claim 4 or 5 further comprising a deletion of up to forty nucleotides positioned 3' to the E1a and E1b and protein IX deletion and a foreign DNA molecule encoding a polyadenylation signal.

7. The recombinant adenovirus expression vector of claims 1 to 6, wherein the adenovirus is a Group C adenovirus selected from a serotype 1, 2, 5 or 6.

8. The recombinant adenovirus expression vector of claim 1, wherein the gene is a DNA molecule up to 2.6 kilobases.

9. The recombinant adenovirus expression vector of claim 6, wherein the gene is a DNA molecule up to 4.5 kilobases.

10. The recombinant adenovirus expression vector of claim 1, wherein the gene encodes a foreign functional protein or a biologically active fragment thereof.

11. The recombinant adenovirus expression vector of claim 10, wherein the gene encodes a foreign functional tumor suppressor protein or a biologically active fragment thereof.

12. The recombinant adenovirus expression vector of claim 1, wherein the gene encodes a suicide protein or functional equivalent thereof.

13. A transformed host cell comprising the recombinant adenovirus expression vector of claim 1 or 10.

14. The transformed host cell of claim 13, wherein the host cell is a procaryotic or eucaryotic cell.

15. A method for transforming a pathologic hyperproliferative mammalian cell comprising contacting the cell with the expression vector of claim 1.

16. A method of treating a pathology in an animal or mammal caused by the absence of a tumor suppressor gene or the presence of a pathologically mutated tumor suppressor gene comprising administering to the  
5 animal or mammal an effective amount of the vector of claim 1 containing a gene encoding a foreign functional protein

0055370 10229  
252207 02585600



having a tumor suppressive function, under suitable conditions.

17. The method of claim 16, wherein the foreign protein is a functional tumor suppressor protein.

18. A method of gene therapy comprising administering to a subject an effective amount of the vector of claim 1.

19. A method of inhibiting the proliferation of a tumor in an animal comprising administering an effective amount of the adenoviral expression vector of claim 1 under suitable conditions to the animal.

20. The method of claim 19, wherein the gene encodes an anti-tumor agent.

21. The method of claim 20, wherein the anti-tumor agent is a tumor suppressor gene.

22. The method of claim 20, wherein the anti-tumor agent is a suicide gene or functional equivalent thereof.

23. The method of claim 21, wherein the tumor is non-small cell lung cancer, small cell lung cancer, hepatocarcinoma, melanoma, retinoblastoma, breast tumor, colorectal carcinoma, leukemia, lymphoma, brain tumor, cervical carcinoma, sarcoma, prostate tumor, bladder tumor, tumor of the reticuloendothelial tissues, Wilm's tumor, astrocytoma, glioblastoma, neuroblastoma, ovarian carcinoma, osteosarcoma, and renal cancer.

24. The method of claim 19, wherein the vector is administered by intra-tumoral injection.

0895370-1089  
2020-05-20

25. A pharmaceutical composition comprising the recombinant adenoviral expression vector of claim 1, 10 or 12.

26. A method for reducing the proliferation of tumor cells in a subject comprising administering under suitable conditions an effective amount of an adenoviral expression vector of claim 12 and an effective amount of a  
5 thymidine kinase metabolite or a functional equivalent thereof.

27. The method of claim 26, wherein the thymidine kinase metabolite is ganciclovir or 6-methoxypurine arabinonucleoside or a functional equivalent thereof.

28. The method of claim 26, wherein the adenoviral expression vector is administered by injection into the tumor mass.

29. The method of claim 26, wherein the tumor cells are hepatocellular carcinoma.

30. The method of claim 29, wherein the adenoviral expression vector is administered directly into the hepatic artery of the subject.

31. A kit for reducing the proliferation of tumor cells comprising the components of the adenoviral expression vector of claim 12, a thymidine kinase metabolite or functional equivalent thereof, pharmaceutical  
5 carriers and instructions for the treatment of hepatocellular carcinoma using the kit components.

0898570-10897


ABSTRACT OF THE DISCLOSURE

This invention provides a recombinant adenovirus expression vector characterized by the partial or total deletion of the adenoviral protein IX DNA and having a gene  
5 encoding a foreign protein or a functional fragment or mutant thereof. Transformed host cells and a method of producing recombinant proteins and gene therapy also are included within the scope of this invention.

Thus, for example, the adenoviral vector of this  
10 invention can contain a foreign gene for the expression of a protein effective in regulating the cell cycle, such as p53, Rb, or mitosin, or in inducing cell death, such as the conditional suicide gene thymidine kinase. (The latter must be used in conjunction with a thymidine kinase  
15 metabolite in order to be effective).

00000000-102897

Figure 1: GENERATION OF A pIX GENE DELETED ADENOVIRUS




(141)  
 (171)  
 (1111)  
 (1511)  
 (1911)  
 (2111)  
 (2711)  
 (3111)  
 (3511)  
 (3911)  
 (4311)  
 (4711)  
 (5111)  
 (5511)  
 (5911)  
 (6311)  
 (6711)  
 (7111)  
 (7511)  
 (7911)  
 (8311)  
 (8711)  
 (9111)  
 (9211)

Single-letter abbreviations


Single-letter abbreviations for the amino acid residues are: A. Ala; C. Cys; D. Asp; E. Glu; F. Phe; G. Gly; H. His; I. Ile; K. Lys; L. Leu; M. Met; N. Asn; P. Pro; Q. Gln; R. Arg; S. Ser; T. Thr; V. Val; W. Trp; and Y. Tyr.

FIGURE 3

123  
243  
343  
408  
428  
438  
458  
478  
498  
518  
538  
558  
578  
598  
618  
638  
658  
678  
698  
718  
738  
758  
778  
798  
818  
838  
858  
878  
898  
918  
938  
958  
978  
998  
1018  
1038  
1058  
1078  
1098  
1118  
1138  
1158  
1178  
1198  
1218  
1238  
1258  
1278  
1298  
1318  
1338  
1358  
1378  
1398  
1418  
1438  
1458  
1478  
1498  
1518  
1538  
1558  
1578  
1598  
1618  
1638  
1658  
1678  
1698  
1718  
1738  
1758  
1778  
1798  
1818  
1838  
1858  
1878  
1898  
1918  
1938  
1958  
1978  
1998  
2018  
2038  
2058  
2078  
2098  
2118  
2138  
2158  
2178  
2198  
2218  
2238  
2258  
2278  
2298  
2318  
2338  
2358  
2378  
2398  
2418  
2438  
2458  
2478  
2498  
2518  
2538  
2558  
2578  
2598  
2618  
2638  
2658  
2678  
2698  
2718  
2738  
2758  
2778  
2798  
2818  
2838  
2858  
2878  
2898  
2918  
2938  
2958  
2978  
2998  
3018  
3038  
3058  
3078  
3098  
3118  
3138  
3158  
3178  
3198  
3218  
3238  
3258  
3278  
3298  
3318  
3338  
3358  
3378  
3398  
3418  
3438  
3458  
3478  
3498  
3518  
3538  
3558  
3578  
3598  
3618  
3638  
3658  
3678  
3698  
3718  
3738  
3758  
3778  
3798  
3818  
3838  
3858  
3878  
3898  
3918  
3938  
3958  
3978  
3998  
4018  
4038  
4058  
4078  
4098  
4118  
4138  
4158  
4178  
4198  
4218  
4238  
4258  
4278  
4298  
4318  
4338  
4358  
4378  
4398  
4418  
4438  
4458  
4478  
4498  
4518  
4538  
4558  
4578  
4598  
4618  
4638  
4658  
4678  
4698  
4718  
4738  
4758  
4778  
4798  
4818  
4838  
4858  
4878  
4898  
4918  
4938  
4958  
4978  
4998  
5018  
5038  
5058  
5078  
5098  
5118  
5138  
5158  
5178  
5198  
5218  
5238  
5258  
5278  
5298  
5318  
5338  
5358  
5378  
5398  
5418  
5438  
5458  
5478  
5498  
5518  
5538  
5558  
5578  
5598  
5618  
5638  
5658  
5678  
5698  
5718  
5738  
5758  
5778  
5798  
5818  
5838  
5858  
5878  
5898  
5918  
5938  
5958  
5978  
5998  
6018  
6038  
6058  
6078  
6098  
6118  
6138  
6158  
6178  
6198  
6218  
6238  
6258  
6278  
6298  
6318  
6338  
6358  
6378  
6398  
6418  
6438  
6458  
6478  
6498  
6518  
6538  
6558  
6578  
6598  
6618  
6638  
6658  
6678  
6698  
6718  
6738  
6758  
6778  
6798  
6818  
6838  
6858  
6878  
6898  
6918  
6938  
6958  
6978  
6998  
7018  
7038  
7058  
7078  
7098  
7118  
7138  
7158  
7178  
7198  
7218  
7238  
7258  
7278  
7298  
7318  
7338  
7358  
7378  
7398  
7418  
7438  
7458  
7478  
7498  
7518  
7538  
7558  
7578  
7598  
7618  
7638  
7658  
7678  
7698  
7718  
7738  
7758  
7778  
7798  
7818  
7838  
7858  
7878  
7898  
7918  
7938  
7958  
7978  
7998  
8018  
8038  
8058  
8078  
8098  
8118  
8138  
8158  
8178  
8198  
8218  
8238  
8258  
8278  
8298  
8318  
8338  
8358  
8378  
8398  
8418  
8438  
8458  
8478  
8498  
8518  
8538  
8558  
8578  
8598  
8618  
8638  
8658  
8678  
8698  
8718  
8738  
8758  
8778  
8798  
8818  
8838  
8858  
8878  
8898  
8918  
8938  
8958  
8978  
8998  
9018  
9038  
9058  
9078  
9098  
9118  
9138  
9158  
9178  
9198  
9218  
9238  
9258  
9278  
9298  
9318  
9338  
9358  
9378  
9398  
9418  
9438  
9458  
9478  
9498  
9518  
9538  
9558  
9578  
9598  
9618  
9638  
9658  
9678  
9698  
9718  
9738  
9758  
9778  
9798  
9818  
9838  
9858  
9878  
9898  
9918  
9938  
9958  
9978  
9998  
10018  
10038  
10058  
10078  
10098  
10118  
10138  
10158  
10178  
10198  
10218  
10238  
10258  
10278  
10298  
10318  
10338  
10358  
10378  
10398  
10418  
10438  
10458  
10478  
10498  
10518  
10538  
10558  
10578  
10598  
10618  
10638  
10658  
10678  
10698  
10718  
10738  
10758  
10778  
10798  
10818  
10838  
10858  
10878  
10898  
10918  
10938  
10958  
10978  
10998  
11018  
11038  
11058  
11078  
11098  
11118  
11138  
11158  
11178  
11198  
11218  
11238  
11258  
11278  
11298  
11318  
11338  
11358  
11378  
11398  
11418  
11438  
11458  
11478  
11498  
11518  
11538  
11558  
11578  
11598  
11618  
11638  
11658  
11678  
11698  
11718  
11738  
11758  
11778  
11798  
11818  
11838  
11858  
11878  
11898  
11918  
11938  
11958  
11978  
11998  
12018  
12038  
12058  
12078  
12098  
12118  
12138  
12158  
12178  
12198  
12218  
12238  
12258  
12278  
12298  
12318  
12338  
12358  
12378  
12398  
12418  
12438  
12458  
12478  
12498  
12518  
12538  
12558  
12578  
12598  
12618  
12638  
12658  
12678  
12698  
12718  
12738  
12758  
12778  
12798  
12818  
12838  
12858  
12878  
12898  
12918  
12938  
12958  
12978  
12998  
13018  
13038  
13058  
13078  
13098  
13118  
13138  
13158  
13178  
13198  
13218  
13238  
13258  
13278  
13298  
13318  
13338  
13358  
13378  
13398  
13418  
13438  
13458  
13478  
13498  
13518  
13538  
13558  
13578  
13598  
13618  
13638  
13658  
13678  
13698  
13718  
13738  
13758  
13778  
13798  
13818  
13838  
13858  
13878  
13898  
13918  
13938  
13958  
13978  
13998  
14018  
14038  
14058  
14078  
14098  
14118  
14138  
14158  
14178  
14198  
14218  
14238  
14258  
14278  
14298  
14318  
14338  
14358  
14378  
14398  
14418  
14438  
14458  
14478  
14498  
14518  
14538  
14558  
14578  
14598  
14618  
14638  
14658  
14678  
14698  
14718  
14738  
14758  
14778  
14798  
14818  
14838  
14858  
14878  
14898  
14918  
14938  
14958  
14978  
14998  
15018  
15038  
15058  
15078  
15098  
15118  
15138  
15158  
15178  
15198  
15218  
15238  
15258  
15278  
15298  
15318  
15338  
15358  
15378  
15398  
15418  
15438  
15458  
15478  
15498  
15518  
15538  
15558  
15578  
15598  
15618  
15638  
15658  
15678  
15698  
15718  
15738  
15758  
15778  
15798  
15818  
15838  
15858  
15878  
15898  
15918  
15938  
15958  
15978  
15998  
16018  
16038  
16058  
16078  
16098  
16118  
16138  
16158  
16178  
16198  
16218  
16238  
16258  
16278  
16298  
16318  
16338  
16358  
16378  
16398  
16418  
16438  
16458  
16478  
16498  
16518  
16538  
16558  
16578  
16598  
16618  
16638  
16658  
16678  
16698  
16718  
16738  
16758  
16778  
16798  
16818  
16838  
16858  
16878  
16898  
16918  
16938  
16958  
16978  
16998  
17018  
17038  
17058  
17078  
17098  
17118  
17138  
17158  
17178  
17198  
17218  
17238  
17258  
17278  
17298  
17318  
17338  
17358  
17378  
17398  
17418  
17438  
17458  
17478  
17498  
17518  
17538  
17558  
17578  
17598  
17618  
17638  
17658  
17678  
17698  
17718  
17738  
17758  
17778  
17798  
17818  
17838  
17858  
17878  
17898  
17918  
17938  
17958  
17978  
17998  
18018  
18038  
18058  
18078  
18098  
18118  
18138  
18158  
18178  
18198  
18218  
18238  
18258  
18278  
18298  
18318  
18338  
18358  
18378  
18398  
18418  
18438  
18458  
18478  
18498  
18518  
18538  
18558  
18578  
18598  
18618  
18638  
18658  
18678  
18698  
18718  
18738  
18758  
18778  
18798  
18818  
18838  
18858  
18878  
18898  
18918  
18938  
18958  
18978  
18998  
19018  
19038  
19058  
19078  
19098  
19118  
19138  
19158  
19178  
19198  
19218  
19238  
19258  
19278  
19298  
19318  
19338  
19358  
19378  
19398  
19418  
19438  
19458  
19478  
19498  
19518  
19538  
19558  
19578  
19598  
19618  
19638  
19658  
19678  
19698  
19718  
19738  
19758  
19778  
19798  
19818  
19838  
19858  
19878  
19898  
19918  
19938  
19958  
19978  
19998  
20018  
20038  
20058  
20078  
20098  
20118  
20138  
20158  
20178  
20198  
20218  
20238  
20258  
20278  
20298  
20318  
20338  
20358  
20378  
20398  
20418  
20438  
20458  
20478  
20498  
20518  
20538  
20558  
20578  
20598  
20618  
20638  
20658  
20678  
20698  
20718  
20738  
20758  
20778  
20798  
20818  
20838  
20858  
20878  
20898  
20918  
20938  
20958  
20978  
20998  
21018  
21038  
21058  
21078  
21098  
21118  
21138  
21158  
21178  
21198  
21218  
21238  
21258  
21278  
21298  
21318  
21338  
21358  
21378  
21398  
21418  
21438  
21458  
21478  
21498  
21518  
21538  
21558  
21578  
21598  
21618  
21638  
21658  
21678  
21698  
21718  
21738  
21758  
21778  
21798  
21818  
21838  
21858  
21878  
21898  
21918  
21938  
21958  
21978  
21998  
22018  
22038  
22058  
22078  
22098  
22118  
22138  
22158  
22178  
22198  
22218  
22238  
22258  
22278  
22298  
22318  
22338  
22358  
22378  
22398  
22418  
22438  
22458  
22478  
22498  
22518  
22538  
22558  
22578  
22598  
22618  
22638  
22658  
22678  
22698  
22718  
22738  
22758  
22778  
22798  
22818  
22838  
22858  
22878  
22898  
22918  
22938  
22958  
22978  
22998  
23018  
23038  
23058  
23078  
23098  
23118  
23138  
23158  
23178  
23198  
23218  
23238  
23258  
23278  
23298  
23318  
23338  
23358  
23378  
23398  
23418  
23438  
23458  
23478  
23498  
23518  
23538  
23558  
23578  
23598  
23618  
23638  
23658  
23678  
23698  
23718  
23738  
23758  
23778  
23798  
23818  
23838  
23858  
23878  
23898  
23918  
23938  
23958  
23978  
23998  
24018  
24038  
24058  
24078  
24098  
24118  
24138  
24158  
24178  
24198  
24218  
24238  
24258  
24278  
24298  
24318  
24338  
24358  
24378  
24398  
24418  
24438  
24458  
24478  
24498  
24518  
24538  
24558  
24578  
24598  
24618  
24638  
24658  
24678  
24698  
24718  
24738  
24758  
24778  
24798  
24818  
24838  
24858  
24878  
24898  
24918  
24938  
24958  
24978  
24998  
25018  
25038  
25058  
25078  
25098  
25118  
25138  
25158  
25178  
25198  
25218  
25238  
25258  
25278  
25298  
25318  
25338  
25358  
25378  
25398  
25418  
25438  
25458  
25478  
25498  
25518  
25538  
25558  
25578  
25598  
25618  
25638  
25658  
25678  
25698  
25718  
25738  
25758  
25778  
25798  
25818  
25838  
25858  
25878  
25898  
25918  
25938  
25958  
25978  
25998  
26018  
26038  
26058  
26078  
26098  
26118  
26138  
26158  
26178  
26198  
26218  
26238  
26258  
26278  
26298  
26318  
26338  
26358  
26378  
26398  
26418  
26438  
26458  
26478  
26498  
26518  
26538  
26558  
26578  
26598  
26618  
26638  
26658  
26678  
26698  
26718  
26738  
26758  
26778  
26798  
26818  
26838  
26858  
26878  
26898  
26918  
26938  
26958  
26978  
26998  
27018  
27038  
27058  
27078  
27098  
27118  
27138  
27158  
27178  
27198  
27218  
27238  
27258  
27278  
27298  
27318  
27338  
27358  
27378  
27398  
27418  
27438  
27458  
27478  
27498  
27518  
27538  
27558  
27578  
27598  
27618  
27638  
27658  
27678  
27698  
27718  
27738  
27758  
27778  
27798  
27818  
27838  
27858  
27878  
27898  
27918  
27938  
27958  
27978  
27998  
28018  
28038  
28058  
28078  
28098  
28118  
28138  
28158  
28178  
28198  
28218  
28238  
28258  
28278  
28298  
28318  
28338  
28358  
28378  
28398  
28418  
28438  
28458  
28478  
28498  
28518  
28538  
28558  
28578  
28598  
28618  
28638  
28658  
28678  
28698  
28718  
28738  
28758  
28778  
28798  
28818  
28838  
28858  
28878  
28898  
28918  
28938  
28958  
28978  
28998  
29018  
29038  
29058  
29078  
29098  
29118  
29138  
29158  
29178  
29198  
29218  
29238  
29258  
29278  
29298  
29318  
29338  
29358  
29378  
29398  
29418  
29438  
29458  
29478  
29498  
29518  
29538  
29558  
29578  
29598  
29618  
29638  
29658  
29678  
29698  
29718  
29738  
29758  
29778  
29798  
29818  
29838  
29858  
29878  
29898  
29918  
29938  
29958  
29978  
29998  
30018  
30038  
30058  
30078  
30098  
30118  
30138  
30158  
30178  
30198  
30218  
30238  
30258  
30278  
30298  
30318  
30338  
30358  
30378  
30398  
30418  
30438  
30458  
30478  
30498  
30518  
30538  
30558  
30578  
30598  
30618  
30638  
30658  
30678  
30698  
30718  
30738  
30758  
30778  
30798  
30818  
30838  
30858  
30878  
30898  
30918  
30938  
30958  
30978  
30998  
31018  
31038  
31058  
31078  
31098  
31118  
31138  
31158  
31178  
31198  
31218  
31238  
31258  
31278  
31298  
31318  
31338  
31358  
31378  
31398  
31418  
31438  
31458  
31478  
31498  
31518  
31538  
31558  
31578  
31598  
31618  
31638  
31658  
31678  
31698  
31718  
31738  
31758  
31778  
31798  
31818  
31838  
31858  
31878  
31898  
31918  
31938  
31958  
31978  
31998  
32018  
32038  
32058  
32078  
32098  
32118  
32138  
32158  
32178  
32198  
32218  
32238  
32258  
32278  
32298  
32318  
32338  
32358  
32378  
32398  
32418  
32438  
32458  
32478  
32498  
32518  
32538  
32558  
32578  
32598  
32618  
32638  
32658  
32678  
32698  
32718  
32738  
32758  
32778  
32798  
32818  
32838  
32858  
32878  
32898  
32918  
32938  
32958  
32978  
32998  
33018  
33038  
33058  
33078  
33098  
33118  
33138  
33158  
33178  
33198  
33218  
33238  
33258  
33278  
33298  
33318  
33338  
33358  
33378  
33398  
33418  
33438  
33458  
33478  
33498  
33518  
33538  
33558  
33578  
33598  
33618  
33638  
33658  
33678  
33698  
33718  
33738  
33758  
33778  
33798  
33818  
33838  
33858  
33878  
33898  
33918  
33938  
33958  
33978  
33998  
34018  
34038  
34058  
34078  
34098  
34118  
34138  
34158  
34178  
34198  
34218  
34238  
34258  
34278  
34298  
34318  
34338  
34358  
34378  
34398  
34418  
34438  
34458  
34478  
34498  
34518  
34538  
34558  
34578  
34598  
34618  
34638  
34658  
34678  
34698  
34718  
34738  
34758  
34778  
34798  
34818  
34838  
34858  
34878  
34898  
34918  
34938  
34958  
34978  
34998  
35018  
35038  
35058  
35078  
35098  
35118  
35138  
35158  
35178  
35198  
35218  
35238  
35258  
35278  
35298  
35318  
35338  
35358  
35378  
35398  
35418  
35438  
35458  
35478  
35498  
35518  
35538  
35558  
35578  
35598  
35618  
35638  
35658  
35678  
35698  
35718  
35738  
35758  
35778  
35798  
35818  
35838  
35858  
35878  
35898  
35918  
35938  
35958  
35978  
35998  
36018  
36038  
36058  
36078  
36098  
36118  
36138  
36158  
36178  
36198  
36218  
36238  
36258  
36278  
36298  
36318  
36338  
36358  
36378  
36398  
36418  
36438  
36458  
36478  
36498  
36518  
36538  
36558  
36578  
36598  
36618  
36638  
36658  
36678  
36698  
36718  
36738  
36758  
36778  
36798  
36818  
36838  
3




**A.**






**B.**







08958570-102897


Media  
Control









A.




B.






268207 02525530


PCR Amplified Fragments









## Genomic Structure of Recombinant Adenoviruses



# Infection with rAANCAT










Figure 15



GREGORY ET AL.  
P-CJ 1192

Figure 16 A and B



DECLARATION AND POWER OF ATTORNEY  
FOR PATENT APPLICATION

As the below-named inventors, we hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled **RECOMBINANT ADENOVIRAL VECTOR AND METHODS OF USE**, the specification of which

\_\_\_\_\_ is attached hereto.  
XXXX was filed on **October 25, 1994**, (Attorney  
Docket No. **P-CJ 1192** ) as Application  
Serial No. **08/328,673**.  
and was amended on (or amended through) \_\_\_\_\_.  
(if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Sec. 1.56(a).

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the

0895370-10399  
"SECRET"

Gregory et al.  
Serial No.: 08/328,673  
Filed: October 25, 1994  
Page 2

national or PCT international filing date of this application:

<u>Application Serial No.</u>	<u>Filing Date</u>	<u>Status</u>
08/233,777	May 19, 1994	Pending
08/142,669	October 25, 1993	Pending

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

We hereby appoint the following attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

CATHRYN CAMPBELL, Registration No. 31,815; CHRISTINE BELLAS, Registration No. 34,122; SUSAN M. PERKINS, Registration No. 36,405; RICHARD J. IMBRA, Registration No. 37,643; PAUL C. STEINHARDT, Registration No. 30,806; GREGORY R. HOOK, Registration No. P38,701; and DAVID SPOLTER, Registration No. 36,933. Direct all telephone calls to Cathryn Campbell at telephone no. (619) 535-9001.

Address all correspondence to:

Cathryn Campbell  
CAMPBELL AND FLORES  
4370 La Jolla Village Drive, Suite 700  
San Diego, California 92122



[illegible]

Inventor's signature: Robert J. Beyer

Post Office Address: 12578 Cavallo Street  
San Diego, California

DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION  
P-CJ 1192  
Serial No.: 08/328,673

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Commissioner of Patents and Trademarks, Washington, D.C. 20231, Attn: Application Processing Division, on February 8, 1995.

By Paul C. Steinhardt  
Paul C. Steinhardt, Reg. No. 30,806  
2/8/95  
Date of Signature

08/328,673

Express Mail Label No. EM197112005US  
Date of Deposit: October 28, 1997


I hereby certify that this correspondence is being deposited  
with the United States Postal Service "Express Mail Post  
Office to Addressee" service under 37 CFR §1.10 on the date  
indicated above and is addressed to:  
Assistant Commissioner for Patents,  
Washington, D.C. 20231,

PATENT

Attorney Docket No. 16930-000921

TOWNSEND and TOWNSEND and CREW LLP

By \_\_\_\_\_  
John P. Borg



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:	)	
	)	
RICHARD J. GREGORY et al.	)	Examiner: Not Assigned
	)	
Application No.: Not Assigned	)	Art Unit: Not Assigned
	)	
Filed: Herewith	)	<u>PRELIMINARY AMENDMENT</u>
	)	
For: ADENOVIRAL PHARMACEUTICAL	)	
VECTOR AND METHODS OF USE	)	

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Prior to examination of the above-referenced  
application, Applicants request entry of the following  
amendments.

IN THE SPECIFICATION:

On page 1, line 4, after "1994", please insert --now  
abandoned--. At line 6, after "1993", please insert --now  
abandoned--.

On page 6, line 12, after "p110<sup>RB</sup>", please insert  
--(SEQ ID NO:8)--. At line 14, after "protein", please insert  
--(SEQ ID NOS:7-8)--.

On page 8, line 21, after "3')", please insert --(SEQ  
ID NO:1)--. At line 22, after "3')", please insert --(SEQ ID  
NO:2)--.

On page 17, line 20, after "Figure 3", please insert  
--(SEQ ID NOS:7-8)--. At line 22, after "Figure 2", please  
insert --(SEQ ID NO:8)--. At line 23, after "Figure 3", please  
insert --(SEQ ID NOS:7-8)--.

18. (Amended) A method of gene therapy comprising administering to a subject an effective amount of [the vector of claim 1] a recombinant adenovirus expression vector comprising a

partial or total deletion of a protein IX DNA sequence and a gene encoding a foreign functional protein having a tumor suppressive function.

19. (Amended) A method of inhibiting the proliferation of a tumor in an animal comprising administering an effective amount of [the adenoviral expression vector of claim 1] a recombinant adenovirus expression vector comprising a partial or total deletion of a protein IX DNA sequence and a gene encoding a foreign functional protein having a tumor suppressive function under suitable conditions to the animal.

26. (Amended) A method for reducing the proliferation of tumor cells in a subject comprising administering under suitable conditions an effective amount of an adenoviral expression vector [of claim 12] comprising a partial or total deletion of a protein IX DNA sequence and a gene encoding a suicide protein or a biologically active fragment thereof and an effective amount of a thymidine kinase metabolite or a functional equivalent thereof.

In claim 31, please delete "12" and insert --26--.

REMARKS

The specification of the present application has been amended to incorporate sequence identifiers, and to incorporate sequence listing pages as required under 37 CFR 1.821. These amendments are formal in nature and present no new matter.

In view of the foregoing, Applicants believe all claims now pending in this application are in condition for early examination and action toward that end is respectfully requested.

RICHARD J. GREGORY et al.  
Application No.: Not Assigned  
Page 4

PATENT

If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at (415) 326-2400.

Respectfully submitted,



Renee A. Fitts  
Reg. No. 35,136

TOWNSEND and TOWNSEND and CREW LLP  
Two Embarcadero Center, 8th Floor  
San Francisco, California 94111-3834  
(650) 326-2400  
Fax (650) 326-2422  
RAF:mc

H:\HOME\RAF\WORK\16930\921\AMEND.PRE

152201" 02525620